

# SpeX Spectroscopy of Unresolved Very Low-Mass Binaries. I. Identification of Seventeen Candidate Binaries Straddling the L Dwarf/T Dwarf Transition

Adam J. Burgasser<sup>1,2,3</sup>, Kelle L. Cruz<sup>3,4</sup>, Michael Cushing<sup>5</sup>, Christopher R. Gelino<sup>6</sup>, Dagny L. Looper<sup>3,5</sup>, Jacqueline K. Faherty<sup>7</sup>, J. Davy Kirkpatrick<sup>3,6</sup>, and I. Neill Reid<sup>3,8</sup>

## ABSTRACT

We report the identification of 17 candidate brown dwarf binaries whose components straddle the L dwarf/T dwarf transition. These sources were culled from a large near-infrared spectral sample of L and T dwarfs observed with the Infrared Telescope Facility SpeX spectrograph. Candidates were selected on the basis of spectral ratios which segregate known (resolved) L dwarf/T dwarf pairs from presumably single sources. Composite templates, constructed by combining 13581 pairs of absolute flux-calibrated spectra, are shown to provide statistically superior fits to the spectra of our seventeen candidates as compared to single templates. Ten of these candidates appear to have secondary components that are significantly brighter than their primaries over the 1.0–1.3  $\mu\text{m}$  band, indicative of rapid condensate depletion at the L dwarf/T dwarf transition. Our results support prior indications of enhanced multiplicity amongst early-type T dwarfs;  $53 \pm 7\%$  of the T0–T4 dwarfs in our spectral sample are found to be either resolved or unresolved (candidate) pairs, although this is consistent with an intrinsic (volume complete) brown dwarf binary fraction of only 15%. If verified, this sample of

---

<sup>1</sup>Center for Astrophysics and Space Science, University of California San Diego, La Jolla, CA 92093, USA; aburgasser@ucsd.edu

<sup>2</sup>Massachusetts Institute of Technology, Kavli Institute for Astrophysics and Space Research, Building 37, Room 664B, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

<sup>3</sup>Visiting Astronomer at the Infrared Telescope Facility, which is operated by the University of Hawaii under Cooperative Agreement no. NCC 5-538 with the National Aeronautics and Space Administration, Science Mission Directorate, Planetary Astronomy Program.

<sup>4</sup>Department of Physics and Astronomy, Hunter College, City University of New York, 695 Park Avenue, New York, NY 10021, USA

<sup>5</sup>Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

<sup>6</sup>Infrared Processing and Analysis Center, California Institute of Technology, Pasadena, CA 91125, USA

<sup>7</sup>Department of Astrophysics, American Museum of Natural History, Central Park West at 79th Street, New York, NY 10034, USA ; Department of Physics and Astronomy, Stony Brook University, Stony Brook, NY 11794-3800, USA

<sup>8</sup>Space Telescope Science Institute, 3700 San Marin Drive, Baltimore, MD 21218, USA

spectral binaries more than doubles the number of known L dwarf/T dwarf transition pairs, enabling a broader exploration of this poorly-understood phase of brown dwarf atmospheric evolution.

*Subject headings:* stars: binaries: general — stars: fundamental parameters — stars: low mass, brown dwarfs

## 1. Introduction

Among the outstanding problems in brown dwarf astrophysics today is the depletion of photospheric condensate clouds at the transition between the two lowest-luminosity spectral classes of low-mass stars and brown dwarfs, the L dwarfs and T dwarfs (Kirkpatrick 2005 and references therein). The L dwarfs, with effective temperatures of  $1300 \lesssim T_{eff} \lesssim 2200$  K (Golimowski et al. 2004), host a variety of mineral condensate species in their photospheres (Lodders 2002), as evident by their very red near-infrared colors, muted H<sub>2</sub>O absorption bands, weak TiO and VO bands (gaseous precursors to condensates) and presence of silicate grain features (e.g., Tsuji et al. 1996; Kirkpatrick et al. 1999; Knapp et al. 2004; Cushing et al. 2006). In contrast, the cooler T dwarfs ( $600 \lesssim T_{eff} \lesssim 1300$  K), which have blue near-infrared colors and strong molecular gas absorption, appear to have relatively condensate-free atmospheres (Marley et al. 1996; Tsuji et al. 1999).

One-dimensional, static cloud models have been able to explain these trends through the confinement of condensates which are vertically bound by a balance of gravitational settling and vertical mixing (e.g., Ackerman & Marley 2001; Tsuji 2002; Burrows et al. 2006). In these models, the cloud layer resides at the photosphere in the L dwarfs, but sinks below the photosphere and out of view in the T dwarfs. While providing a robust, conceptual framework for understanding cloud evolution at the L dwarf/T dwarf transition, these models have nevertheless failed to explain the surprisingly narrow range of effective temperatures ( $\Delta T_{eff} \approx 200\text{--}400$  K) and luminosities ( $\Delta \log L_{bol}/L_{\odot} \approx 0.3$  dex) that encompass this transition (Kirkpatrick et al. 2000; Golimowski et al. 2004; Burgasser 2007a). More remarkably, surface fluxes in the  $1.0\text{--}1.3\ \mu\text{m}$  region are observed to *increase* amongst early-type T dwarfs, the so-called “*J*-band bump” (Dahn et al. 2002; Tinney et al. 2003). Careful tuning of cloud parameters (e.g, settling efficiency, cloud coverage or cloud top temperature) can reproduce the characteristics of the L dwarf/T dwarf transition, but in an arguably ad-hoc manner (e.g., Burgasser et al. 2002b; Tsuji 2005; Cushing et al. 2008).

Binaries with L dwarf and T dwarf components have helped to clarify this transition by providing coeval, co-spatial laboratories for detailed atmospheric investigations. Resolved photometry for a handful of these systems (Burgasser et al. 2006c; Liu et al. 2006;Looper et al. 2008b) have confirmed the *J*-band bump to be an intrinsic aspect of brown dwarf evolution, rather than variations in surface gravity, metallicity or unresolved multiplicity in heterogeneous samples (Tsuji & Nakajima 2003; Burrows et al. 2006). L/T binaries have also enabled better characterization of absolute magnitude/spectral type relations across the L dwarf/T dwarf transition, confirming the narrow

luminosity range over which this transition occurs. Interpreted as a short timescale for the removal of condensates from brown dwarf photospheres ( $\sim 500$  Myr; Burgasser 2007a), this result also explains the relative rarity of early-type T dwarfs (Metchev et al. 2008) and their high rate of multiplicity (nearly twice that of other brown dwarfs in magnitude-limited samples; Burgasser et al. 2006c; Liu et al. 2006).

Two factors conspire to hamper resolved imaging studies of L dwarf/T dwarf pairs: their low space density and corresponding greater distances from the Sun, and the inherently small physical separations of brown dwarf multiples in general ( $\sim 98\%$  have  $a \lesssim 20$  AU; e.g., Allen 2007). Fortunately, the distinct and highly-structured near-infrared spectral energy distributions of L and T dwarfs makes it possible to both identify and characterize *unresolved* pairs from combined-light near-infrared spectroscopy, so-called spectral binaries. Indeed, one of the first L/T binaries found, 2MASS J0518-2828<sup>1</sup>, was originally identified by its peculiar near-infrared spectrum (Cruz et al. 2004), and subsequently confirmed via high angular resolution imaging with the *Hubble Space Telescope* (*HST*; Burgasser et al. 2006c). A more recent spectral binary discovery, 2MASS J0320-0446 (Burgasser et al. 2008a), was independently identified as an M8.5 + T5 spectroscopic binary with an orbital period of 0.67 yr (Blake et al. 2008). This case illustrates the increased potential of finding very closely-separated systems among unresolved binaries, systems that are more amenable to astrometric and/or spectroscopic orbital mass measurements that provide critical tests of brown dwarf evolutionary theories (e.g., Zapatero Osorio et al. 2004; Liu et al. 2008; Dupuy et al. 2009). Tight binaries also have a higher probability of eclipsing (e.g., Stassun et al. 2006) and hence the potential to yield radius measurements to constrain as-yet untested structural models of evolved brown dwarfs (i.e., ages  $\gtrsim 1$  Gyr; Chabrier et al. 2009). Unresolved pairs can be identified at greater distances, enabling the construction of larger and more statistically robust samples; and provide a means of measuring the true binary fraction of brown dwarfs currently uncertain by a factor of 2-3 due to separation selection biases (e.g., Lodieu et al. 2007; Joergens 2008).

In this article, we report the identification of seventeen unresolved, L dwarf/T dwarf binary candidates selected from a large sample of low-resolution near-infrared spectra of late-type dwarfs. In Section 2 we summarize the construction and basic characteristics of the sample, including new observations of L and T dwarfs. In Section 3 we describe our method of selecting L dwarf/T dwarf binary candidates through the use of spectral indices. In Section 4 we detail our spectral template analysis technique, define metrics to assess the probability of unresolved multiplicity, and determine component parameters. In Section 5 we present the results of our spectral analysis, examining each candidate in detail. In Section 6 we discuss our results in the context of the absolute magnitude

---

<sup>1</sup>Names of sources listed in the text are shortened to [PREFIX] Jhhmm±ddmm, where [PREFIX] is the original discovery catalog: 2MASS (Two Micron All Sky Survey; Skrutskie et al. 2006), SDSS (Sloan Digital Sky Survey; York et al. 2000), DENIS (Deep Near-Infrared Survey of the Southern Sky; Epchtein et al. 1997) or IPMS (Infrared Proper Motion Survey; Artigau et al. 2006), and hhmm and ±ddmm are the sexagesimal Right Ascension (hours and minutes) and declination (degrees and arcminutes) at equinox J2000. Full source names and coordinates are given in Tables 2, 3 and 7; additional source data can be found at the DwarfArchives site, <http://dwarfarchives.org>.

relations for L and T dwarfs, the frequency of multiples across the L dwarf/T dwarf transition, and the process of cloud depletion at this transition. The major conclusions of this study are summarized in Section 7. As Sections 2 through 4 are devoted to our selection and analysis techniques, readers interested primarily in the binary candidates can skip to Section 5 without loss of context.

## 2. SpeX Spectral Sample

Our analysis is based on a large sample of low-resolution, near-infrared spectra of L and T dwarfs obtained with SpeX (Rayner et al. 2003), a low-resolution, 0.8–5.4  $\mu\text{m}$  spectrograph mounted on the 3m NASA Infrared Telescope Facility. The spectra were obtained using the prism-dispersed mode of SpeX, which provides 0.75–2.5  $\mu\text{m}$  continuous wavelength coverage in a single order, with dispersion of 20–30  $\text{\AA pixel}^{-1}$  and an average resolution  $\lambda/\Delta\lambda \approx 120$  for the 0".5 slit (some sources were also observed with the 0".8 slit). The sample is neither complete nor volume-limited; it constitutes a representative subset of ultracool dwarfs in the local disk population, based on ongoing SpeX spectroscopic follow-up by ourselves and other researchers of all known L and T dwarfs down to an effective magnitude limit of  $K \approx 16$ , including nearly all known L and T dwarfs within 20 parsecs of the Sun. Most of these sources observed are compiled on the Dwarf Archives site<sup>2</sup>; the SpeX spectra examined here are available at the SpeX Prism Spectral Libraries<sup>3</sup>.

### 2.1. New Observations

Ninety-nine L and T dwarfs were observed by the authors over several runs spanning 2004 September through 2008 September, as summarized in Table 1. These data were acquired in a variety of weather (clear to partly cloudy) and seeing conditions (0".3–1".2), but were uniformly observed using the 0".5 or 0".8 slit aligned at parallactic angle and generally at low airmass ( $\sec z \lesssim 2$ ). Each source was observed in multiple exposures, dithering in an ABBA pattern along the slit, with total integration times of 240–1200 s depending on source brightness and weather conditions. Nearby A0 V stars were observed for flux calibration and telluric absorption correction, and internal flat field and argon arc lamps were observed with each flux standard for pixel response and wavelength calibration. All data were reduced using the SpeXtool reduction package (Vacca et al. 2003; Cushing et al. 2004) using standard settings, as described in detail in Burgasser (2007b).

---

<sup>2</sup><http://dwarfarchives.org>.

<sup>3</sup><http://www.browndwarfs.org/spexprism>.

## 2.2. Data from the Literature

In addition to our new observations, we compiled from the literature SpeX observations of 134 L and T dwarfs with published optical and/or near-infrared spectral types spanning L3 to T8 and with data having median signal-to-noise ratios of 20 or greater over the 0.9–2.4  $\mu\text{m}$  window. Those data<sup>4</sup> were acquired and reduced following similar procedures as described above. Combining the new and previous observations results in a sample of 253 spectra for 233 L3–T8 dwarfs.

## 2.3. Initial Template Sample

In order to identify and characterize unresolved binaries, we required a “clean” sample of spectral templates. An initial sample was constructed by excluding known binaries (Burgasser et al. 2007b and references therein) young cluster objects (e.g., Muench et al. 2007), and sources specifically noted to have peculiar spectra associated with low surface gravities, subsolar metallicities or unusual cloud properties (e.g., Lucas & Roche 2000; Burgasser et al. 2003a; Knapp et al. 2004; Chiu et al. 2006; Cruz et al. 2007, 2009;Looper et al. 2008b). This left us with a sample of 189 spectra of 178 sources, whose properties are summarized in Table 2.

Figure 1 shows the distribution of published spectral types for these spectra, based on optical data for L3–L8 dwarfs or near-infrared data for L9–T8 dwarfs and those L dwarfs without optical classifications. The distribution is relatively flat, albeit with fewer T0–T3 dwarfs and T8 dwarfs reflecting the rarity of these sources in current search samples. The literature classifications are based primarily on the Kirkpatrick et al. (1999), Geballe et al. (2002) or Burgasser et al. (2006b) schemes. However, many L dwarfs have never been typed in the near-infrared, and those with near-infrared types have been classified following different schemes (e.g., Reid et al. 2001a; Testi et al. 2001; McLean et al. 2003; Nakajima et al. 2004; Burgasser 2007a). To place the sample on a self-consistent, homogeneous basis, we calculated near-infrared spectral types for each source directly from the SpeX data, using the spectral indices H<sub>2</sub>O-J, CH<sub>4</sub>-J, H<sub>2</sub>O-H, CH<sub>4</sub>-H, H<sub>2</sub>O-K and CH<sub>4</sub>-K defined in Burgasser et al. (2006b) and index/spectral types relations given in Burgasser (2007a). As in the latter study, the set of indices used to classify each source is selected based on an initial (in this case, published) spectral type estimate. The H<sub>2</sub>O-J, H<sub>2</sub>O-H and H<sub>2</sub>O-K indices are combined for types L8 and earlier and the H<sub>2</sub>O-J, CH<sub>4</sub>-J, H<sub>2</sub>O-H, CH<sub>4</sub>-H and CH<sub>4</sub>-K indices are used for later types. As there can be significant differences between published and near-infrared spectral types, particularly for the L dwarfs (e.g., Geballe et al. 2002), we repeated the classification process three times for each source, using each iteration’s classification to determine the indices used for the next iteration. The spectral type adopted for each source is the mean of its individual index

---

<sup>4</sup>Data were drawn from Burgasser et al. (2004, 2006a,b, 2007a, 2008a,b); Cruz et al. (2004); Burgasser & McElwain (2006); Chiu et al. (2006); McElwain & Burgasser (2006); Reid et al. (2006); Burgasser (2007a,b,c); Liebert & Burgasser (2007);Looper et al. (2007, 2008b); Luhman et al. (2007); Siegler et al. (2007) and Sheppard & Cushing (2009).

types.

Table 2 lists the index-based types, with uncertainties (standard deviation of index types) given for those sources with considerable scatter (default classification uncertainty is  $\pm 0.5$  subtypes). Figure 1 shows the distribution of these classifications, which is not significantly different from the literature distribution. Figure 2 compares the SpeX and literature classifications against each other. Over the entire template sample, the standard deviation between classifications is 0.9 subtypes, although average deviations are generally larger among the L dwarfs ( $\sigma = 1.1$  subtypes) than the T dwarfs ( $\sigma = 0.5$  subtypes). In most cases, the discrepancies are between optical and near-infrared types for L dwarfs. Only eleven sources had SpeX classifications that differed by more than two subtypes from the published classifications, one of which is a candidate binary (2MASS J1711+2232; see Section 3.2). For completeness, we used both published and index-based spectral types in our analysis.

### 3. Identification of Binary Candidates

#### 3.1. Spectral Peculiarities in Known Binary Systems

Prior identification of unresolved L dwarf/T dwarf binaries from SpeX data has been based largely on the presence of peculiar spectral features, such as the sharp dip at  $1.6\ \mu\text{m}$  observed in the spectra of 2MASS J0320-0446, SDSS J0805+4812 (Burgasser 2007b) and Kelu-1A (Stumpf et al. 2008) attributed to overlapping FeH and CH<sub>4</sub> absorption features from unresolved L dwarf and T dwarf components. However, such specific features need not arise in all L/T binary combinations; see for example Figure 4 in Burgasser (2007a).

In an effort to make the selection process more robust, we examined the near-infrared spectra of six resolved L/T binaries (Table 3): 2MASS J0518-2828, SDSS J0423-0414 (Burgasser et al. 2005b), SDSS J1021-0304 (Burgasser et al. 2006c), 2MASS J1404+3159 (Looper et al. 2008a), SDSS J1534+1615 (Liu et al. 2006), and 2MASS J2252-1730 (Reid et al. 2006). Figure 3 compares the combined light spectrum of each of these systems to its closest match in our template library (see Section 4.2). Spectral deviations between the binaries and templates range from pronounced (in the case of 2MASS J0518-2828) to subtle (in the case of SDSS J1021-0304), but several common trends stand out:

1. The  $1.6\ \mu\text{m}$  CH<sub>4</sub> absorption band is typically stronger relative to  $2.2\ \mu\text{m}$  band in the binaries, the former occasionally appearing in the absence of the latter (e.g., 2MASS J0518-2828);
2. The  $1.27\ \mu\text{m}$  flux peak is more pronounced and H<sub>2</sub>O and CH<sub>4</sub> absorption deeper at  $1.1\ \mu\text{m}$  in the binaries (e.g., SDSS J1534+1615);
3. The  $2.1\ \mu\text{m}$  flux peak is slightly shifted toward the blue in the binaries (e.g., SDSS J1021-0304); and

4. The overall near-infrared spectral energy distribution is generally bluer in the binaries (e.g., 2MASS J2252-1730).

These traits suggest qualitative means of differentiating unresolved binaries from single sources, and systems such as 2MASS J0518-2828 readily stand out in direct spectral comparisons. However, comparative methods implicitly assume that the templates themselves are “normal” single spectra, an assumption we cannot make given that we are searching for unresolved binaries within the same sample. To overcome this ambiguity, we used a suite of spectral indices to identify sources whose overall spectral properties are similar to known binaries and therefore indicative of unresolved multiplicity.

### 3.2. Spectral Index Selection

Eight spectral indices, defined in Table 4, were measured for each spectrum in our full spectral sample (templates and known binaries/peculiar sources). These include the six classification indices used in Section 2.2, the  $K/J$  flux peak ratio defined in Burgasser et al. (2006b), and an additional index sampling the  $1.6\ \mu\text{m}$  feature noted above. We compared all eight indices against each other for every source in our sample, and examined trends in individual indices and index ratios as a function of spectral type using both literature and index-based classifications.

From these combinations, we identified six pairings that best segregated known L dwarf/T dwarf binaries from the bulk of the spectral sample, as shown in Figure 4. These pairings reflect the peculiar traits outlined above, in particular the strength of the  $H$ -band  $\text{CH}_4$  feature ( $\text{CH}_4\text{-K}$  versus  $\text{CH}_4\text{-H}$  and  $\text{H-dip}$  versus  $\text{H}_2\text{O-H}$ ), the unusual brightness of the  $J$ -band peak ( $\text{H}_2\text{O-K}$  versus  $\text{H}_2\text{O-J}$ ,  $\text{H}_2\text{O-J/H}_2\text{O-H}$  and  $\text{H}_2\text{O-J/CH}_4\text{-K}$  versus spectral type), and blue near-infrared colors ( $\text{K/J}$  versus  $\text{CH}_4\text{-H}$ ). In these comparison spaces, we identified regions (Table 5) which clearly segregated known binaries, but were conservative enough so as not to over-select unusual spectra that presumably arise from non-multiplicity effects (e.g., variations in surface gravity, metallicity or cloud properties). The criteria also allowed us to retain enough non-binary-like late-L/early-T dwarf spectra for our spectral template analysis (Section 4). These criteria clearly provide an incomplete selection of unresolved binaries, as they do not select several known binary systems with either weak signatures of  $\text{CH}_4$  absorption (e.g., 2MASS J0850+1057 and 2MASS J1728+3948; Reid et al. 2001b; Gizis et al. 2003) or systems with identical components (e.g., SDSS J0926+584; Burgasser et al. 2006c). However, as a first attempt at identifying unresolved pairs, we chose to maximize the reliability of our candidate selection over sample completeness.

In all, 46 sources from our full SpeX sample of 253 spectra satisfied at least one selection criterion. These include nine known binaries and all six sources listed in Table 3. We refined our selection by requiring binary candidates to satisfy at least two criteria. We also segregated “strong” candidates (satisfying at least three criteria) from “weak” candidates (satisfying only two criteria), a distinction intended to test the robustness of the spectral index criteria. These 20 sources are

listed in Table 7, and constitute our initial candidate pool. All have near-infrared classifications from SpeX data spanning L8.5–T3; several have been previously noted in the literature as having peculiar or highly uncertain spectral types. We discuss each candidate in detail in Section 5.

## 4. Spectral Template Analysis

### 4.1. Flux Calibration

The final test of the binary nature of our candidates and characterization of their components is a comparison of their spectra against synthetic composites generated from the template spectra. The templates, purged of binary candidates (reducing the sample to 170 spectra of 161 sources), were interpolated onto a common wavelength scale and set on an absolute flux scale (in  $F_\lambda$  flux units) using the MKO<sup>5</sup>  $M_K$ /spectral type relations of Liu et al. (2006). We considered the two  $M_K$  relations defined in that study, one constructed by rejecting known binaries (hereafter, the “bright” relation) and one constructed from rejecting known and candidate binaries (hereafter, the “faint” relation; see Figure 5 in Liu et al. 2006). These two relations encompass our current best constraints on absolute magnitude trends across the L dwarf/T dwarf transition, despite diverging by nearly  $\sim 1$  mag over spectral types L8–T5. Absolute magnitudes were assigned according to either literature or index-based spectral types, resulting in four independent flux scalings for the templates. As our baseline calibration set, we adopted the faint calibration applied to literature classifications (see Section 6.1).

### 4.2. Comparison to Single and Composites

For each candidate, we first determined best matches to individual template spectra following a procedure similar to that outlined in Burgasser et al. (2008a). All spectra were initially normalized to the maximum flux in the 1.2–1.3  $\mu\text{m}$  region. We then computed a weighted chi-square statistic between each candidate ( $C[\lambda]$ ) and template ( $T[\lambda]$ ) spectrum:

$$\chi^2 \equiv \sum_{\{\lambda\}} w[\lambda] \left[ \frac{C[\lambda] - \alpha T[\lambda]}{\sigma_c[\lambda]} \right]^2 \quad (1)$$

(see Cushing et al. 2008). Here,  $w[\lambda]$  is a vector of weights satisfying  $\sum_{\{\lambda\}} w[\lambda] = 1$ ,  $\alpha$  is a scaling factor that minimizes  $\chi^2$  (see Equation 2 in Cushing et al. 2008),  $\sigma_c[\lambda]$  is the noise spectrum for the candidate, and the sum is performed over the wavelength ranges  $\{\lambda\} = 0.95\text{--}1.35 \mu\text{m}$ ,  $1.45\text{--}1.8 \mu\text{m}$  and  $2.0\text{--}2.35 \mu\text{m}$  in order to avoid regions of strong telluric absorption. We adopted the same weighting scheme used in Cushing et al. (2008), with each pixel weighted by its spectral width (i.e.,  $w_i \propto \Delta\lambda_i$ ).

---

<sup>5</sup>Mauna Kea Observatory filter system; see Tokunaga et al. (2002) and Simons & Tokunaga (2002).



The candidate spectra were then compared to a larger set of synthetic composites, constructed by combining all possible pairs of flux-calibrated templates for which one source was of equal or later spectral type (using either literature or index-based classifications). The resulting 13581 composites were also normalized and compared to the candidate spectra using the same  $\chi^2$  statistic as Equation 1. We note that because we are comparing the candidate spectra against a finite sample of spectra of distinct sources, each of which are modulated by some degree of noise, our expectation is not to achieve  $\chi^2 \approx 1$  for our best-fit cases; indeed, this expectation is not even realized in spectral model fits (see Figure 4 in Cushing et al. 2008). What matters is whether a binary template provides a significantly improved fit, as discussed below.

### 4.3. Assessing Fit Quality and Parameter Estimation

The number of composites used to fit the candidate spectra vastly outnumbers the number of single templates, so we are almost assured of finding a better fit (lower  $\chi^2$ ) with the former. It is therefore necessary to assess the statistical significance of the fit improvement in order to rule out the null hypothesis; i.e., that the candidate is not a binary. We used the one-sided F-test for this purpose, using as our distribution statistic the ratio

$$\eta_{SB} \equiv \frac{\min(\{\chi_{single}^2\})/\nu_{single}}{\min(\{\chi_{composite}^2\})/\nu_{binary}} \quad (2)$$

where  $\nu$  is the degrees of freedom in each fit. For both single and composite fits, one might initially assume that  $\nu_{single} = \nu_{composite} = \nu$  is the number of data points used in the fit ( $N = 296$ ), minus one to account for the relative scaling between data and template spectra. However, individual weighting of the spectral points implies that they do not contribute equally to the total  $\chi^2$  value. We therefore define the effective number of data points,

$$N_{eff} \equiv \frac{1}{\max(\{w\})} \sum_{i=1}^N w_i \quad (3)$$

which reduces to  $N$  for  $w_i = \text{constant}$ . For our weighting scheme,  $N_{eff} = 254$  and hence  $\nu = 253$ . To rule out our null hypothesis at the 99% confidence level (CL), we required  $\eta_{SB} > 1.34$  as our final selection criteria.

Typically, multiple single or composite templates yield similar  $\chi^2$  values for a given candidate. We therefore computed mean values and uncertainties for the component parameters (i.e., spectral types and relative brightnesses) using a weighting scheme based on the F-distribution. In effect, each fit’s parameter set was weighted by the likelihood that that fit is equivalent to the best fit:

$$W_i \propto 1 - F(\eta_{i0} \mid \nu, \nu). \quad (4)$$

Here,  $\eta_{i0} \equiv \chi_{i0}^2 / \min([\chi^2])$  is the ratio of chi-square residuals between the best-fit template and the  $i^{th}$  template, and  $F(\eta_{i0} \mid \nu, \nu)$  is the F-distribution probability distribution function. Parameter

means ( $\bar{p}$ ) and uncertainties ( $\sigma_p$ ) were then computed as

$$\bar{p} \equiv \frac{\sum_i W_i p_i}{\sum_i W_i} \quad (5)$$

and

$$\sigma_p^2 = \frac{\sum_i W_i (p_i - \bar{p})^2}{\sum_i W_i}, \quad (6)$$

where the sums are over all fits.

#### 4.4. Known L/T Transition Binaries

Before proceeding to examine our binary candidates, we first assessed how well our index selection and spectral fitting procedures identified and reproduced the properties of the known binary systems listed in Table 3. The best-fit templates for these sources are shown in Figure 5, while Table 5 summarizes the component parameters based on the faint calibration scale. For each system, best-fit composites provided a statistically significant better match (CL > 99%) to the combined light spectrum than the best-fit single template. Visual inspection of these fits also indicates clear improvement in the cases of 2MASS J2252-1730, SDSS J0423-0414, 2MASS J0518-2828 and 2MASS J1404+3159. The fits for SDSS J1021-0304 and SDSS J1534+1615 are more subtle improvements, but notably reproduce the blue-shifted peak at  $K$ -band in the spectrum of the former and the strong 1.1  $\mu\text{m}$  H<sub>2</sub>O/CH<sub>4</sub> band in the spectrum of the latter.

Resolved photometric measurements have been made for these systems, so we examined how well our technique reproduces reported relative magnitudes. For the composites, synthetic MKO  $JHK$  and  $HST/NICMOS$   $F110W$  and  $F170M$  photometry were computed directly from the flux-calibrated SpeX data by integrating these and a Kurucz model spectrum of Vega with the respective filter profiles (see Cushing et al. 2005). Mean relative magnitudes and their uncertainties were calculated as in Equations 5 and 6, and are also listed in Table 6. In all six cases, synthetic photometry is consistent with measurements to within  $2\sigma$ , although uncertainties in the former are as high as 0.8 mag. For the two sources with the most reliable photometry, SDSS J0423-0414 and 2MASS J1404+3159, the agreement is within  $1.5\sigma$ .

Finally, while none of the known binaries listed in Table 3 have reported resolved spectroscopy, the inferred component types are generally consistent with estimated types from the literature ( $\pm 1$  subtype) with the exceptions of 2MASS J0518-2828, for which we infer later-type components (L7.5+T5 versus L6+T4:), and DENIS J2252-1730, for which we infer an earlier primary and later secondary (L4.5+T4.5 versus L6+T2). However, reported types for the former are highly uncertain (Cruz et al. 2004; Burgasser et al. 2006c) while our spectral fit to the latter is superior to that presented in Reid et al. (2006).

In summary, we find that our method is capable of robustly identifying and characterizing known L dwarf/T dwarf transition pairs, providing a measure of confidence in our candidate binary

selection and component characterization.

## 5. Results

Single and composite template fits for each of the candidate binaries and for all four component flux calibrations are summarized in Tables 8 and 9. These include best-fit and mean component types and relative MKO *JHK* and *HST*/NICMOS *F110W* and *F170M* magnitudes. The best fits based on the faint flux calibration are shown in Figures 6 and 7. In the following sections we discuss each candidate in detail.

### 5.1. Strong Candidates

#### 5.1.1. *SDSS J024749.90-163112.6*

Originally identified in the SDSS by Chiu et al. (2006), SDSS J0247-1631 is classified  $T2\pm1.5$  in the near-infrared, consistent with our index-based classification of  $T2.5\pm1$ . The uncertainty in both reflects large scatter in spectral indices, with stronger absorption in the  $1.1\ \mu\text{m}$   $\text{H}_2\text{O}/\text{CH}_4$  band relative to  $1.6$  and  $2.2\ \mu\text{m}$   $\text{CH}_4$  bands. SDSS J0247-1631 satisfied all six spectral index selection criteria, and the best-fit composite is a clear improvement over the single template fit and statistically significant ( $\text{CL} > 99\%$ ). Average primary and secondary spectral types are inferred to be  $T0\pm0.2$  and  $T7\pm0.3$ , the latter being the latest-typed secondary in our sample. The composition of this system indicates a secondary that is fainter than the primary across the near-infrared band ( $\Delta J = 0.68\pm0.10$ ). The primary component of the best-fit composite, SDSS J1520+3546, has discrepant literature ( $T0\pm1$ ) and index-based (L7.5) near-infrared classifications, due in part to weak  $\text{H}_2\text{O}$  absorption at  $1.1\ \mu\text{m}$  and  $1.3\ \mu\text{m}$ ; and its near-infrared color ( $2\text{MASS } J - K_s = 1.54\pm0.08$ ) is somewhat blue for either type. Faherty et al. (2009) have found that unusually blue L dwarfs have a large tangential velocity ( $V_{\text{tan}}$ ) distribution, indicative of older ages, higher surface gravities and possibly subsolar metallicities. SDSS J0247-1631 itself has a relatively large  $V_{\text{tan}} = 46\pm7\ \text{km s}^{-1}$ , compared to a mean of  $26\pm19\ \text{km s}^{-1}$  for L0–L9 dwarfs (Faherty et al. 2009). These characteristics suggest that SDSS J0247-1631 may be an older and/or slightly metal-poor pair of brown dwarfs. No high angular resolution imaging observations of this source have been reported to date.

#### 5.1.2. *SDSS J035104.37+481046.8*

Chiu et al. (2006) classified SDSS J0351+4810 a  $T1\pm1$  in the near-infrared, similar to our SpeX classification ( $T1.5\pm1$ ). Again, the uncertainties arise from spectral peculiarities, in this case unusually weak  $\text{CH}_4$  absorption at  $2.2\ \mu\text{m}$ . SDSS J0351+4810 satisfied all six selection criteria,

and the best-fit composite is a clear improvement over the best-fit single template and statistically significant. Average primary and secondary spectral types are  $L6.5 \pm 0.7$  and  $T5 \pm 0.7$ , with a secondary that may be slightly brighter than the primary at the  $1.05 \mu\text{m}$  and  $1.27 \mu\text{m}$  spectral flux peaks ( $\Delta J = 0.31 \pm 0.31$ ). The primary in the best-fit composite, 2MASS J0103+1935 (L6, 2MASS  $J - K_s = 2.14 \pm 0.10$ ), has a fairly steep red spectral slope, indicative of either low surface gravity (i.e., youth) or unusually thick condensate clouds (e.g., Looper et al. 2008b). On the other hand, the best-fit secondary has a normal color for its spectral type. This combination suggests a system with an unusually cloudy primary and normal secondary. No high angular resolution imaging observations of SDSS J0351+4810 have been reported to date.

### 5.1.3. SDSS J103931.35+325625.5

Chiu et al. (2006) classified SDSS J1039+3256 a T1 in the near-infrared (our SpeX classification is T1.5), and its spectrum satisfied all six selection criteria. The best-fit composite is a clear improvement over the best-fit single template and statistically significant. Average primary and secondary spectral types are  $L7 \pm 0.2$  and  $T4 \pm 0.2$ , and the primary appears to be brighter than the secondary at all wavelengths ( $\Delta J = 0.26 \pm 0.09$ ). The components of the best-fit composite appear to be fairly normal for their respective spectral types. No high angular resolution imaging observations of SDSS J1039+3256 have been reported to date.

### 5.1.4. 2MASS J11061197+2754225

Looper et al. (2007) classified 2MASS J1106+2754 a T2.5 in the near-infrared; we classified the same SpeX spectrum T2 using spectral indices. Satisfying three selection criteria, 2MASS J1106+2754 is subtly but significantly better fit by composites with average component types of  $T0 \pm 0.2$  and  $T4.5 \pm 0.2$ . These types are similar to those inferred for 2MASS J1404+3159, and predict a secondary that is substantially brighter at the  $1.05 \mu\text{m}$  and  $1.27 \mu\text{m}$  spectral flux peaks ( $\Delta J = -0.37 \pm 0.06$ ). Again, the best-fit primary is SDSS J1520+3546, a somewhat peculiar blue late-L/early-T dwarf.

2MASS J1106+2754 is one of only two strong candidates imaged at high angular resolution, in this case by Looper et al. (2008a) using the Keck AO system.  $K_s$ -band images show only a single point source with a point spread function full width at half maximum of 68 mas. Based on the relative magnitudes inferred from this analysis ( $\Delta K = 1.16 \pm 0.09$  for the faint calibration), this observation rules out a resolved binary system with projected separation  $\gtrsim 1.5$  AU at the time of the observation, assuming a distance of 22 pc (Looper et al. 2007). The null imaging result indicates one of three possibilities: 2MASS J1106+2754 may be a single source, a binary observed close to line-of-sight alignment (e.g., Kelu-1; Martin et al. 1999; Liu & Leggett 2005; Gelino et al. 2006), or a tightly-separated system. The excellent fit of this source’s spectrum to composites supports one of the two latter possibilities, and 1.5 AU is not a particularly stringent constraint on the separation

given that several brown dwarf multiple systems are known to have even smaller separations (e.g., Basri & Martín 1999; Stassun et al. 2006; Joergens & Müller 2007). Nevertheless, second-epoch imaging and/or spectroscopic monitoring are required to verify the binary nature of this source.

#### 5.1.5. *2MASS J13243559+6358284*

Discovered independently by Looper et al. (2007) and Metchev et al. (2008), 2MASS J1324+6358 was classified as a peculiar T2 in the former study based on its unusually red near-infrared spectral energy distribution (our SpeX classification is T2±1). 2MASS J1324+6358 is also a red outlier in optical, near-infrared and mid-infrared colors (Looper et al. 2007; Metchev et al. 2008; Faherty et al. 2009). Looper et al. specifically examined the possibility that this source is an unresolved binary, finding a good match to an L9 + T2 composite. We also find this source to be a likely binary, with statistically significant better fits to composites with mean component types of L8±0.2 and T3.5±0.2. The secondary of this system appears to be brighter than the primary at the 1.05  $\mu\text{m}$  and 1.27  $\mu\text{m}$  flux peaks ( $\Delta J = -0.05 \pm 0.06$ ). The primary component of the best-fit composite, 2MASS J1043+2225 (2MASS  $J - K_s = 1.97 \pm 0.08$ ), has been noted as one of the reddest optically-classified L8 dwarfs known (Cruz et al. 2007), while the secondary has normal colors for its spectral type. Like SDSS J0351+4810, this combination suggests a system with an unusually cloudy primary and cloud-free T dwarf secondary. No high angular resolution imaging observations of 2MASS J1324+6358 have been reported to date.

#### 5.1.6. *SDSS J141530.05+572428.7*

Chiu et al. (2006) classified SDSS J1415+5724 a T3±1 (same as our SpeX classification), with an uncertainty driven by weak 2.2  $\mu\text{m}$  CH<sub>4</sub> absorption relative to the pronounced 1.6  $\mu\text{m}$  band. Both Chiu et al. (2006) and Faherty et al. (2009) note this source as being unusually red for its spectral type, and the former attempted (unsuccessfully) to reproduce its spectrum as a binary. SDSS J1415+5724 satisfied all six selection criteria, and composites (average types L8±0.5 and T5±0.3) provide visually obvious and statistically significant better fits to its spectrum. The secondary component appears to be brighter than the primary at the 1.05  $\mu\text{m}$  and 1.27  $\mu\text{m}$  spectral peaks ( $\Delta J = -0.13 \pm 0.20$ ), and both components in the best-fit composite have normal colors and spectral energy distributions for their respective spectral types. No high angular resolution imaging observations of SDSS J1415+5724 have been reported to date.

#### 5.1.7. *SDSS J143553.25+112948.6*

Chiu et al. (2006) classified SDSS J1435+1129 a T2±1 in the near-infrared (consistent with our T2.5±1 SpeX classification), with the uncertainty driven by weak 2.2  $\mu\text{m}$  CH<sub>4</sub> absorption relative to

1.1  $\mu\text{m}$  and 1.6  $\mu\text{m}$  bands. Its spectrum satisfied all six selection criteria, and the best-fit composite is a visually obvious better match than the best-fit single template, reproducing in particular the broad  $K$ -band peak and relatively strong 1.6  $\mu\text{m}$   $\text{CH}_4$  absorption. The average component types,  $\text{L}7.5 \pm 0.4$  and  $\text{T}6 \pm 0.3$ , indicate a secondary that is slightly fainter than the primary throughout the near-infrared ( $\Delta J = 0.41 \pm 0.12$ ). The components of the best-fitting composite appear to have normal colors and spectral energy distributions for their respective spectral types. No high angular resolution imaging observations of SDSS J1435+1129 have been reported to date.

#### 5.1.8. SDSS J143945.86+304220.6

Chiu et al. (2006) classified SDSS J1439+3042 a T2.5, identical to our index-based classification. Its spectrum satisfied only four selection criteria, but the best-fit composite is a statistically significantly (albeit subtle) better match than the best-fit single template, providing a better fit to the strong 1.6  $\mu\text{m}$   $\text{CH}_4$  band and blue-shifted  $K$ -band peak. Average component types of  $\text{T}1 \pm 0.2$  and  $\text{T}5 \pm 0.6$  indicate a secondary component that may be slightly brighter than the primary at the 1.27  $\mu\text{m}$  spectral flux peak ( $\Delta J = 0.06 \pm 0.24$ ). The best-fit binary components have normal colors and spectral energy distributions for their respective types. No high angular resolution imaging observations of SDSS J1439+3042 have been reported to date.

#### 5.1.9. SDSS J151114.66+060742.9

Chiu et al. (2006) classified SDSS J1511+0607 a  $\text{T}0 \pm 2$  in the near-infrared (our SpeX type is  $\text{T}0.5 \pm 2$ ), with the large uncertainty arising from substantial scatter in index subtypes (L7–T2) on the Geballe et al. (2002) scheme. Its spectrum is clearly peculiar, with strong  $\text{CH}_4$  absorption at 1.1  $\mu\text{m}$  and 1.6  $\mu\text{m}$ , but very weak absorption at 2.2  $\mu\text{m}$ ; these features are similar in nature to the resolved binary 2MASS J0518-2828. The spectrum of SDSS J1511+0607 satisfied all six selection criteria, and the best-fit composite is a visually obvious and statistically significant better match than the best-fit single template. Average component types,  $\text{L}5.5 \pm 0.8$  and  $\text{T}5 \pm 0.4$ , indicate a secondary component that is fainter than the primary across the near-infrared band ( $\Delta J = 0.54 \pm 0.32$ ). The primary component of the best-fit composite, SDSS J0624-4521 (L5, 2MASS  $J - K_s = 1.88 \pm 0.04$ ) is somewhat red for its spectral type, while the secondary has normal colors. Hence, this may be another system with an unusually cloudy primary but normal secondary. No high angular resolution imaging observations of SDSS J1511+0607 have been reported to date.

#### 5.1.10. SDSS J151603.03+025928.9

Knapp et al. (2004) classified SDSS J1516+0259 a  $\text{T}0 \pm 1.5$  on the near-infrared scheme of Geballe et al. (2002); our SpeX classification is  $\text{L}9.5 \pm 1$ . The uncertainties in both arise from weak

CH<sub>4</sub> absorption at 2.2  $\mu\text{m}$  compared to the 1.1  $\mu\text{m}$  and 1.6  $\mu\text{m}$  bands. The spectrum of this source satisfied four of the spectral index criteria, and the best-fit composite is a significantly better match than the best-fit single template, 2MASS J0328+2302, which may itself be a unresolved binary (see Section 6.1). The average component types inferred for SDSS J1516+0259, L7.5 $\pm$ 1.1 and T2.5 $\pm$ 2.2, are rather poorly constrained, as are the relative brightnesses ( $\Delta J = 0.30 \pm 0.65$ ). Notably, the best-fit composite combines a somewhat red L7 (2MASS J0318-3421, 2MASS  $J - K_s = 2.06 \pm 0.07$ ) and a fairly normal T2.5, again suggesting a system with a cloudy primary but normal secondary. No high angular resolution imaging observations of SDSS J1516+0259 have been reported to date.

#### 5.1.11. 2MASS J1711457+223204

2MASS J1711+2232 has been studied extensively in the literature, originally identified in the 2MASS survey and classified L6.5 in the optical with no detectable lithium absorption or H $\alpha$  emission (Kirkpatrick et al. 2000). We derive a later but poorly constrained near-infrared classification of L9 $\pm$ 3 based on strong H<sub>2</sub>O absorption and an indication of weak CH<sub>4</sub> absorption at 1.6  $\mu\text{m}$ . 2MASS J1711+2232 was also examined in the near-infrared by Nakajima et al. (2004), who noted the presence of CH<sub>4</sub> at 2.2  $\mu\text{m}$  but not at 1.6  $\mu\text{m}$ ; our SpeX spectrum in fact shows the opposite, although the  $K$ -band data have somewhat lower signal-to-noise. Tsuji et al. (2004) commented on the unusual 1.6  $\mu\text{m}$  spectral morphology of this source, suggesting that it may be due to absorption from FeH and possibly some other unidentified species.

That other species is likely to be CH<sub>4</sub> from an unresolved T dwarf companion, as the unusual feature at 1.6  $\mu\text{m}$  seen in the SpeX spectrum of 2MASS J1711+2232 (which satisfied all six selection criteria) is similar to those noted in the spectra of 2MASS J0320-0446, SDSS J0805+4812 and Kelu-1A. The best-fit composite is a significantly better match than our best-fit single template, reproducing the 1.6  $\mu\text{m}$  feature in detail. Mean component types of L5 $\pm$ 0.4 and T5.5 $\pm$ 1.2 are similar to those inferred for SDSS J0805+4812 (L4.5 $\pm$ 0.7 and T5 $\pm$ 0.6; Burgasser 2007b). The secondary is considerably fainter than the primary across the near-infrared ( $\Delta J = 0.92 \pm 0.32$ ,  $\Delta K = 3.05 \pm 0.60$ ).

2MASS J1711+2232 has been imaged at high angular resolution with *HST*/WFPC2 (Gizis et al. 2003) and reported to be unresolved in both F814W ( $\lambda_c = 0.79 \mu\text{m}$ ) and F1042M ( $\lambda_c = 1.02 \mu\text{m}$ ) bands. However, the sensitivity of these observations were insufficient to detect the T dwarf component; the primary/combined source was marginally detected at F1042M, where the much fainter secondary ( $\Delta F1042M \approx 2-3$ ) would have been brightest. Hence, in the absence of sufficiently sensitive near-infrared *HST* or AO observations, we cannot determine conclusively whether this system is a tight (unresolveable) binary). 2MASS J1711+2232 is the only binary candidate in our sample with a parallactic distance measurement ( $d = 30 \pm 4$  pc; Vrba et al. 2004). Its component absolute magnitudes are discussed further in Section 6.1.

### 5.1.12. 2MASS J21392676+0220226

2MASS J2139+0220 was classified T0 by Reid et al. (2008) based on its red optical spectrum, and T1.5 in the near-infrared by Burgasser et al. (2006b) based on the SpeX spectrum examined here (indices yield a  $T2.5 \pm 1$  spectral type). Its near-infrared color is somewhat red for its spectral type: 2MASS  $J - K_s = 1.68 \pm 0.07$  as compared to  $\langle J - K_s \rangle = 1.31$  for T1 dwarfs and  $\langle J - K_s \rangle = 1.02$  for T2 dwarfs (Faherty et al. 2009). The spectrum of 2MASS J2139+0220 satisfied only three selection criteria, but this source nevertheless appears to be a strong binary candidate as its best-fit composite is a significant improvement over the best-fit single template. However, the fit is not perfect, failing to reproduce the weak  $1.6 \mu\text{m}$   $\text{CH}_4$  band relative to strong  $1.1 \mu\text{m}$  and  $2.2 \mu\text{m}$  absorptions. The inferred average component types,  $L8.5 \pm 0.7$  and  $T3.5 \pm 1.0$ , are also somewhat poorly constrained. Given its unusual pattern of  $\text{CH}_4$  features even for an unresolved binary, the components of 2MASS J2139+0220 may themselves have unusual properties. Its best-fitting primary, the  $L9 \pm 1$  SDSS J0830+4828, has a large tangential velocity ( $V_{tan} = 79 \pm 4 \text{ km s}^{-1}$ ; Faherty et al. 2009) although its absolute brightness and infrared colors appear to be normal (Geballe et al. 2002; Vrba et al. 2004; Leggett et al. 2007). The inferred composition of this system indicates a secondary that is substantially brighter than the primary at the  $1.05 \mu\text{m}$  and  $1.27 \mu\text{m}$  spectral peaks ( $\Delta J = -0.14 \pm 0.21$ ). No high angular resolution imaging observations of 2MASS J2139+0220 have been reported to date.

## 5.2. Weak Candidates

### 5.2.1. SDSS J011912.22+240331.6

Chiu et al. (2006) classified SDSS J0119+2403 a T2 in the near-infrared (similar to our index-based T2.5 classification), and its spectrum exhibits a slightly enhanced  $J$ -band peak and stronger  $1.6 \mu\text{m}$   $\text{CH}_4$  absorption relative to the T2 spectral standard, SDSS J1254-0122. The best-fit composite is a subtle but significant improvement over the best-fit single template, with inferred component types of  $T0 \pm 0.7$  and  $T4 \pm 0.4$ , similar to those of the resolved binary 2MASS J1404-3159. Like that source, the secondary of SDSS J0119+2403 appears to be substantially brighter than the primary at the  $1.05 \mu\text{m}$ ,  $1.27 \mu\text{m}$  and possibly  $1.6 \mu\text{m}$  spectral peaks ( $\Delta J = -0.42 \pm 0.19$ ). The best-fit primary, SDSS J0858+3256 (T1, MKO  $J - K = 1.61 \pm 0.04$ ) is noted to be a red outlier for its spectral type with a large tangential velocity ( $V_{tan} = 66 \pm 3 \text{ km s}^{-1}$ ; Faherty et al. 2009), while the best-fit secondary is an unexceptional T4 dwarf. Hence, this system may again be composed of an unusually cloudy primary and cloud-free secondary. No high angular resolution imaging observations of SDSS J0119+2403 have been reported to date.



### 5.2.2. *SDSS J075840.33+324723.4*

Knapp et al. (2004) classified SDSS J0758+3247 a T2±1 on the Geballe et al. (2002) scheme, with an uncertainty driven by strong H<sub>2</sub>O/CH<sub>4</sub> absorption at 1.1 μm (our SpeX classification is T2.5). This source is in fact a near-clone to SDSS J1254-0122, and our binary templates actually provide significantly *worse* fits (higher  $\chi^2$  values) than this best-fit single template. While SDSS J1254-0122 also satisfied one of our binary selection criteria, it is unresolved in *HST* imaging observations (Burgasser et al. 2006c) so we cannot exclude the possibility that both sources are single brown dwarfs. No high angular resolution imaging observations of SDSS J0758+3247 have been reported to date.

### 5.2.3. *SDSS J090900.73+652527.2*

Chiu et al. (2006) classified SDSS J0909+6525 a T1.5 in the near-infrared, identical to our SpeX classification, and its spectrum is fairly similar to that of the bright T2.5 IPMS J0136+0933. The MKO  $J - K = 0.62 \pm 0.04$  color of SDSS J0909+6525 is somewhat blue for its spectral type, and Faherty et al. (2009) specifically note it as a blue outlier with a normal tangential velocity ( $V_{tan} \approx 28 \text{ km s}^{-1}$ ). The best-fit composite is a subtle but significantly better match to the spectrum of SDSS J0909+6525, indicating average component types of T1.5±0.5 and T2.5±0.3 and a secondary that is somewhat brighter than the primary at the 1.05 μm and 1.27 μm spectral peaks ( $\Delta J = -0.12 \pm 0.10$ ). The best-fitting composite components have normal colors and spectral energy distributions for their respective spectral types. No high angular resolution imaging observations of SDSS J0909+6525 have been reported to date.

### 5.2.4. *2MASS J09490860-1545485*

Tinney et al. (2005) classified 2MASS J0949-1545 a T1±1 in the near-infrared (our SpeX classification is T1.5), the uncertainty arising from strong 1.6 μm CH<sub>4</sub> absorption but weak 1.1 μm H<sub>2</sub>O/CH<sub>4</sub> absorption as compared to spectral standards. The spectrum of this source is also similar to that of IPMS J0136+0933, but the best-fit composite is a significantly better match with components similar to SDSS J0909+6525, T1±0.2 and T2±0.2. Again, the secondary is inferred to be somewhat brighter than the primary at the 1.05 μm and 1.27 μm spectral peaks ( $\Delta J = -0.07 \pm 0.05$ ) despite having comparable spectral types. The best-fit composite components are normal for their respective classifications. No high angular resolution imaging observations of 2MASS J0949-1545 have been reported to date.

### 5.2.5. *SDSS J120602.51+281328.7*

Chiu et al. (2006) classified SDSS J1206+2813 a T3 in the near-infrared (identical to our SpeX classification), and it exhibits somewhat blue near-infrared colors for its spectral type, MKO  $J-K = 0.13 \pm 0.04$  versus  $\langle J-K \rangle = 0.56 \pm 0.22$  for other T2.5–T3.5 dwarfs (Chiu et al. 2006). Its spectrum is nearly identical to that of the T3.5 SDSS J1750+1759, and while our best-fit composite provides a slight reduction in  $\chi^2$ , at 92% confidence it falls just short of our significance threshold. It is possible that SDSS J1206+2813 fails the binary significance criteria because SDSS J1750+1759 is itself an unresolved binary, as suggested by its apparent overluminosity compared to other early-type T dwarfs (Vrba et al. 2004; Liu et al. 2006). However, SDSS J1750+1759 was unresolved in *HST* images (Burgasser et al. 2006c) and it did not satisfy any of our spectral index selection criteria. Hence, we cannot rule out that both sources are single brown dwarfs. No high angular resolution images of SDSS J1206+2813 have been reported to date.

### 5.2.6. *SDSS J120747.17+024424.8*

Hawley et al. (2002) classified SDSS J1207+0244 an L8 based on red optical data, while Knapp et al. (2004) classified it T0 in the near-infrared (identical to our SpeX classification). The spectrum of SDSS 1207+0244 is very similar to that of the resolved binary SDSS J0423-0414, and as a consequence it was chosen to replace the latter as the T0 spectral standard in the classification scheme of Burgasser et al. (2006b). However, it appears that this source may also be a binary system, as the best-fit composite provides a significantly better match to its spectrum. The mean component types,  $L6.5 \pm 0.7$  and  $T2.5 \pm 0.5$  are similar to those inferred for SDSS J0423-0414 (Table 6), the secondary being fainter than the primary throughout the near-infrared ( $\Delta J = 0.48 \pm 0.28$ ). The components of the best-fitting composite are normal, although the T3 2MASS J1209-1004 satisfied one of our selection criteria. The possibility that SDSS J1207+0244 is a binary system with very different component types is potentially problematic for the currently-defined T dwarf classification scheme. High angular resolution images of this source have yet to be reported.

### 5.2.7. *SDSS J151643.01+305344.4*

Chiu et al. (2006) classified SDSS J1516+3053 a  $T0.5 \pm 1$  while our SpeX classification is  $T1.5 \pm 2$ . The uncertainties in both arise from this source’s unusually red spectral energy distribution and anomalously strong H<sub>2</sub>O absorption bands. Leggett et al. (2007) also note this source as being unusually red in mid-infrared colors, and comparison to model colors suggest the presence of strong vertical mixing and thick condensate clouds in its atmosphere (Ackerman & Marley 2001; Saumon & Marley 2008). Chiu et al. (2006) attempted to fit their spectral data to a binary model, but did not find an adequate match. Our analysis concurs with that result, as the best-fit composite is only a marginal improvement over the best-fit single template (87% confidence level), and notably

fails to reproduce the strong  $\text{H}_2\text{O}/\text{CH}_4$  absorption feature at  $1.1\ \mu\text{m}$ . We therefore cannot rule out the possibility that this source is a single brown dwarf with unusual atmospheric properties. No high angular resolution imaging observations of SDSS J1516+3053 have been reported to date.

#### 5.2.8. SDSS J205235.31-160929.8

Chiu et al. (2006) classified SDSS J2052-1609 a T1 $\pm$ 1 (our SpeX classification is T0 $\pm$ 1), the uncertainty arising in part from strong absorption at  $1.1\ \mu\text{m}$  as compared to weak  $\text{H}_2\text{O}$  and  $\text{CH}_4$  features at longer wavelengths. Despite satisfying only two selection criteria, this source appears to be a strong binary candidate, as the best-fit composite is a subtle but significantly better match to the data than the best-fit single template. Average component types of L7.5 $\pm$ 0.6 and T2 $\pm$ 0.2 are similar to those of SDSS J0423-0414, although in this case the secondary is inferred to be slightly brighter than its primary at the  $1.05\ \mu\text{m}$  and  $1.27\ \mu\text{m}$  spectral peaks ( $\Delta J = 0.04\pm 0.18$ ). The components of the best-fit composite, SDSS J1155-0559 (L7; 2MASS  $J - K_s = 1.54\pm 0.11$ ) and 2MASS J1122-3512 (T2; 2MASS  $J - K_s = 0.64\pm 0.08$ ), both have unusually blue near-infrared colors for their respective spectral types ( $\langle J - K_s \rangle = 1.81$  and  $1.02$  for L7 and T2 dwarfs, respectively; Faherty et al. 2009), suggesting that SDSS J2052-1609 may be an older or slightly metal-poor system. Indeed, its large proper motion ( $\mu = 0''.483\pm 0''.022\ \text{yr}^{-1}$ ; J. Faherty, in preparation) implies a fairly large tangential velocity ( $V_{\text{tan}} = 48\pm 8\ \text{km s}^{-1}$ ), even without accounting for unresolved multiplicity in the photometric distance estimate ( $21\pm 3\ \text{pc}$ ). No high angular resolution imaging observations of this source have been reported to date.

#### 5.2.9. Summary

Based on our analysis, we find that 17 of the 20 binary candidates identified by our spectral index selection criteria have a high probability of being (as yet) unresolved binaries, as indicated by the statistically significant better fits provided by composites. These include all of our strong candidates but only 63% (5/8) of our weak candidates, suggesting that at least three index selection criteria are preferred to identify robust binary candidates. The inferred properties of these systems are summarized in Table 10. The twelve systems with component types between L7 and T5 more than doubles the number of known binaries whose components span the L dwarf/T dwarf transition,<sup>6</sup> although we emphasize that these systems remain candidates pending verification via high angular resolution imaging and/or high resolution spectroscopic monitoring.

---

<sup>6</sup>Known or suspected L/T transition binaries include the six sources listed in Table 3 plus 2MASS J0850+1057 (Reid et al. 2001b; Dahn et al. 2002), 2MASS J0920+3517 (Reid et al. 2001b; Burgasser et al. 2006b), 2MASS J1728+3948 (Gizis et al. 2003),  $\epsilon$  Indi BC (McCaughrean et al. 2004), and Gliese 337CD (Burgasser et al. 2005a).

## 6. Discussion

### 6.1. Absolute Magnitudes Across the L Dwarf/T Dwarf Transition

As discussed in Section 1, there remains considerable uncertainty in absolute magnitude trends across the L dwarf/T dwarf transition due to the presence of unresolved multiples, gravity and metallicity effects, and ultimately the behavior of condensate clouds as they disperse out of the photosphere. With a large number of candidate binaries in our sample whose components straddle this transition, we considered whether it was possible to use these systems to better constrain these trends.

We first considered the quality of fits provided by the composites for the two flux calibration relations of Liu et al. (2006), which differ by nearly a full magnitude between types L7 and T5. For the seventeen sources identified as promising binary candidates, we find that for 65% (11/17) of these the bright relation provides a lower  $\chi^2$  value for the best-fit composite (the same fraction is derived whether literature or index-based classifications are used). However, for only one case—SDSS J1207+0244—does the bright relation provide a statistically significant better fit; i.e., CL >99% in the ratio of minimum  $\chi^2$  values between the best-fit composites for the bright and faint relations. On the other hand, for SDSS J1039+3256 and 2MASS J1324+6358 the bright relation provides a statistically significant *worse* fit (CL <1%). Figure 8 compares the best-fit composites based on both faint and bright flux calibrations for these three sources, and indeed the lower  $\chi^2$  fits are clear improvements. All three of these systems also contain a T2–T4 secondary, within the spectral type range where the faint and bright relations of Liu et al. (2006) differ the most. Given the inconsistency in which relation provides better fits for our candidate binaries, we cannot definitely select one over the other on this analysis alone.

A more direct assessment can be made by focusing on individual brown dwarfs (singles and binary components) with parallax distance measurements. Figure 9 displays absolute MKO  $J$  magnitudes<sup>7</sup> for 58 L and T dwarfs with measured parallaxes and absolute magnitude errors  $\leq 0.3$  mag. Eighteen of these sources are known binaries; we include component magnitudes of the only candidate binary in our sample with a parallax measurement, 2MASS J1711+2232, along with measured component magnitudes for SDSS J0423-0414, SDSS J1021-0304 and  $\epsilon$  Indi BC (McCaughrean et al. 2004). Nearly all of the unresolved sources and binary components follow the faint  $M_J$  relation up to spectral type  $\sim$ T2; only the L9.5 2MASS J0328+2302 stands out as clearly overluminous in this range, with  $M_J$  magnitudes comparable to combined light photometry for Gliese 337CD. Between types T2 and T5, there are only two unresolved sources with parallax measurements, SDSS J1750+1759 (T3.5) and 2MASS J0559-1404 (T4.5), both of which lie well above the faint relation; indeed, the latter lies  $\sim 0.6$  mag above the *bright*  $M_J$  relation. This could indicate a

---

<sup>7</sup>Photometry shown in Figure 9 is based either on direct measurements (Geballe et al. 2002; Leggett et al. 2002; Knapp et al. 2004; Chiu et al. 2006) or synthetic filter corrections from 2MASS photometry as computed directly from the SpeX data; see Section 4.4.

dramatic brightening over this spectral type range, as suggested by the component brightnesses of the resolved binary 2MASS J1404+3159 (Looper et al. 2008a) and our unresolved candidate SDSS J0119+2403 (Section 5.2.1). On the other hand, SDSS J1750+1759 and 2MASS J0559-1404 have both been suggested as unresolved multiples (Golimowski et al. 2004; Liu et al. 2006; Stephens et al. 2009), although neither satisfy any of our selection criteria. We therefore cannot rule out the possibility that they are young or otherwise peculiar sources.

The paucity of absolute magnitude measurements in the T2–T5 range makes it impossible to determine at this point whether absolute  $J$  magnitudes remain relatively flat across the entire L/T transition (a “ $J$ -band plateau”; Liu et al. 2006), favoring of modest flux reversal and more gradual cloud depletion; or undergo a sharp increase over types T2–T5 (a “ $J$ -band spike”; Looper et al. 2008a), requiring very rapid cloud depletion. Indeed, neither of these scenarios may encompass the behavior of all brown dwarfs. Increasing the number of T2–T5 dwarfs with parallax measurements, and obtaining resolved photometry of the candidates presented here, are both necessary if we are to properly constrain trends and distributions of absolute magnitudes across the L/T transition.

## 6.2. The Frequency of Brown Dwarf Binaries

Higher rates of multiplicity amongst early-type T dwarfs have been observed in resolved imaging studies (Burgasser et al. 2006c), although statistics in these samples remain poor due to small numbers (Goldman et al. 2008). As our candidate sample includes sources that might be undetectable in high resolution imaging, it is relevant to assess how their inclusion in multiplicity statistics modifies brown dwarf binary fraction trends across the L/T transition, and what constraint can be made on the intrinsic binary fraction.

As discussed in Section 2, our spectral sample is neither complete nor volume-limited, and selection biases are uncertain given that sources were identified from different search programs using different color and magnitude selection criteria. However, as a first approach we made the simplifying assumption that the sample is roughly magnitude-limited, and estimated the fraction of L8–T6 dwarfs with SpeX data that are either resolved or candidate binaries. These numbers are summarized in Table 11, and indicate the same high multiplicity rate amongst early-type T dwarfs as found in resolved imaging studies. For T0–T4 dwarfs, we deduce a binary fraction of  $53 \pm 7\%$  (23/43 sources), peaking above 60% for T0–T2 dwarfs alone. Over half of the early-type T dwarfs in this spectral sample appear to be multiples, compared to  $\sim 30\%$  in resolved imaging samples (Burgasser 2007a). Moreover, the fraction inferred here is likely to be a lower limit, given that we are unable to identify unresolved binaries with identical components or combinations that fall outside our spectral index selection criteria (see Section 3.2).

Such a high *apparent* rate of multiplicity does not necessary translate into a high *intrinsic* (i.e., volume-limited) rate, however. As discussed in Burgasser (2007a), the enhanced binary fraction of early-type T dwarfs arises predominantly from the small change in luminosity across the L dwarf/T

dwarf transition, which induces both a sharp minimum in the luminosity function of brown dwarfs at type T0 (roughly three times rarer in number than early- and mid-type L dwarfs) and makes late-type L and early-type T dwarfs comparably bright in the near-infrared. Combined with the preferential selection of binaries in magnitude-limited samples, these effects collude to amplify the apparent multiplicity rate of early-type T dwarfs. To infer the intrinsic rate, we reproduced the Monte Carlo population simulations of Burgasser (2007a), using the baseline parameters defined in that study and assuming a magnitude-limited sample. We find that an intrinsic rate of only 15% can produce an apparent binary fraction of 56% for T0–T4 dwarfs in a magnitude-limited sample, comparable to the fraction estimated above. This intrinsic rate is in fact on the low end of current estimates that attempt to correct for unresolved multiplicity (e.g., Basri & Reiners 2006; Lodieu et al. 2007; Joergens 2008), and remains consistent with a brown dwarf multiplicity fraction that is significantly lower than those of more massive stars (e.g., Duquennoy & Mayor 1991; Fischer & Marcy 1992). We reiterate, however, that our result constitutes a lower limit, and a robust measure requires detailed modeling of selection effects (e.g., component spectral type sensitivity range, mass ratio sensitivity as a function of age) that are beyond the scope of this paper.

### 6.3. Flux Reversals and Cloud Evolution

The physical feature that ties both absolute magnitude trends and enhanced binary fractions across the L/T transition is the “sudden” depletion of condensate clouds from the photosphere. This is particularly evident in the increased surface fluxes at the  $1.05\ \mu\text{m}$  and  $1.27\ \mu\text{m}$  flux peaks where cloud particles are a dominant source of opacity (Ackerman & Marley 2001). Amongst the binary candidates identified here, 59% (10/17) of the entire sample and 82% (9/11) of candidates with L7–T5 components show similar flux reversals in the  $J$ -band region. While these reversals are dictated in part by our adopted flux calibration (which are, however, tied to the  $K$ -band flux; see Section 4.1), the quality of the fits coupled with observed absolute magnitude/spectral type trends (Section 6.1) suggest that these flux reversals are real.

With a relatively large sample of possible L dwarf/T dwarf pairs, we can also assess how these flux reversals vary between systems with similar spectral compositions. Consider the candidates SDSS J0119+2403 and SDSS J1106+2754, which have component types of T0+T4 and T0+T4.5 and relative brightnesses of  $\Delta J = -0.42 \pm 0.19$  and  $-0.37 \pm 0.06$ , respectively. Examination of the best-fit composites for these sources (Figures 6 and 7) reveals that the former has a far more pronounced flux reversal than the latter, with the secondary of SDSS J0119+2403 being potentially brighter even at the  $1.6\ \mu\text{m}$  flux peak. The best-fit primary of this system was inferred to be an unusually cloudy source, which translates in greater opacity in the  $J$ -band region. In contrast, the primary of 2MASS J1106+2754 was inferred to be slightly blue for its spectral type, which could be attributed to thinner condensate clouds (e.g., Knapp et al. 2004; Burgasser et al. 2008b). If clouds are the main driver for the unusual colors of the primary components, and their secondaries are

roughly equivalent, the different flux reversals can be tied directly to variations in cloud properties. Specifically, an initially cloudy brown dwarf should lose more opacity in the  $J$ -band region, and experience a greater increase in brightness, as it evolves across the L/T transition as compared to a brown dwarf that initially has thinner clouds. This correlation between initial cloud content and degree of flux reversal suggests that absolute magnitude trends may have more spread across the L/T transition than can be discerned in the currently sparse dataset; spread has also been suggested to arise from age/surface gravity effects (e.g., Metchev & Hillenbrand 2006; Mohanty et al. 2007) and metallicity variations (e.g., Burrows et al. 2006). Again, resolved studies of confirmed binaries in our sample would provide insight into how atmospheric properties modulate the removal of condensate clouds.

#### 6.4. Limitations of this Study

Despite the large number of potential L dwarf/T dwarf pairs uncovered in this investigation, there are clearly limitations as pertaining to the completeness of the sample and robustness of the component characterizations. We reiterate that the 17 candidates listed in Table 10 may not represent the full complement of potential binaries in our spectral sample. Our selection criteria were specifically chosen to be conservative so as to not eliminate too many spectral templates, particularly over the L8–T5 spectral type range. As such, we failed to select known binary systems with early-type T dwarf classifications (Section 3.2) and a number of suspected binary pairs such as 2MASS J0328+2302. We are also unable to select systems with similar spectral component types, and are likely biased against higher-order multiples such as Kelu 1 (Stumpf et al. 2008) and DENIS-P J020529.0-115925 (Bouy et al. 2005). The retention of unresolved binaries in our template sample may also skew the statistical significance of our candidates. For instance, our rejected candidate SDSS J1206+2813, whose spectrum is a near clone to that of SDSS J1750+1759, may prove to be a binary after all should the latter be determined as such. On the other hand, incomplete sampling of single L dwarf/T dwarf transition objects in our spectral template library may cause us to overselect candidates, a possibility for the poorly-fit candidate 2MASS J2139+0220. It is in this vein that we encourage follow-up high resolution imaging (or second epoch imaging in the cases of 2MASS J1106+2754 and 2MASS J1711+2232) and spectroscopy of the candidates listed in Table 10 to both verify their binary nature and better characterize their components.

### 7. Summary

We have identified 17 candidate brown dwarf binaries whose components straddle the L dwarf/T dwarf transition. Their unresolved multiplicity is inferred from similarities in near-infrared spectral indices to known binary systems, and statistically significant better fits to near-infrared spectra by composite templates as compared to single templates. Ten of these systems appear to have secondaries that are brighter than their primaries over the 1–1.3  $\mu\text{m}$  region, by up to  $\sim 0.4$  mag,

consistent with a nonequilibrium depletion of condensate clouds across the L dwarf/T dwarf transition. This is despite the fact that our analysis of these systems is based largely on the faint absolute magnitude relations of Liu et al. (2006). We cannot rule out an even more pronounced *J*-band flux reversal for the sparsely sampled T3–T5 dwarfs. We find that  $53 \pm 7\%$  of the T0–T4 dwarfs in our SpeX spectral sample are resolved or unresolved (candidate) binaries, a rate that is consistent with an intrinsic brown dwarf binary fraction of at least 15%, assuming the sample is magnitude-limited. We find some evidence of flux reversal variations between similarly-classified pairs that may arise from cloud effects, although these must be verified through more accurate characterization of the components. While the sample of candidate binaries presented here does require verification through follow-up high resolution imaging and/or spectroscopic monitoring, it constitutes a promising collection of coeval laboratories for understanding the atmospheric processes that drive the still poorly-understood L dwarf/T dwarf transition.

The authors would like to thank telescope operators Bill Golisch, Dave Griep and Paul Sears, and instrument specialist John Rayner, for their assistance during our many IRTF runs. This publication makes use of data from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center, and funded by the National Aeronautics and Space Administration and the National Science Foundation. 2MASS data were obtained from the NASA/IPAC Infrared Science Archive, which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This research has also made use of the SIMBAD database, operated at CDS, Strasbourg, France; the M, L, and T dwarf compendium housed at DwarfArchives.org and maintained by Chris Gelino, Davy Kirkpatrick, and Adam Burgasser; the SpeX Prism Spectral Libraries, maintained by Adam Burgasser at <http://www.browndwarfs.org/spexprism>; and the VLM Binaries Archive maintained by Nick Siegler at <http://www.vlmbinaries.org>. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

Facilities: IRTF (SpeX)






Fig. 1.— Distribution of spectral types for the template sample based on classifications from the literature (left) and calculated from the SpeX data (right). The former are based on optical types for L3–L8 dwarfs (where available) and near-infrared types for L9–T8 dwarfs. The distributions are roughly consistent with each other and relatively flat for the small number of sources available per subtype.




Fig. 2.— Comparison of spectral types from the literature and those calculated directly from the SpeX data. Overall, classifications are consistent to within 0.9 subtypes, although there is considerably more scatter among the L dwarfs ( $\sigma = 1.1$ ) than the T dwarfs ( $\sigma = 0.5$ )




Fig. 3.— Comparison of known (resolved) binary spectra (Table 3; black lines) to best-fit single templates (red lines). Observed data (in  $F_\lambda$  flux units) have been normalized to the peak flux in the 1.0–1.3  $\mu\text{m}$  range; templates are relatively normalized to minimize  $\chi^2$  deviations. Source names for both the binary and best-fit template are listed in top right corner of each panel, with both published classifications and index-based classifications (in parentheses) indicated. Also shown are the noise spectra (uncertainty in flux as a function of wavelength; grey lines) for the binaries.




Fig. 4.— Spectral index selection criteria for binary candidates. Top four panels compare pairs of index values— $H_2O-K$  versus  $H_2O-J$ ,  $CH_4-K$  versus  $CH_4-H$ ,  $K/J$  versus  $CH_4-H$  and  $H\text{-dip}$  versus  $H_2O-H$ —with estimated spectral types for each abscissa indicated at top. Bottom two panels compare index ratios  $H_2O-J/H_2O-H$  and  $H_2O-J/CH_4-K$  to near-infrared spectral type as derived from the SpeX data. In all panels, templates are indicated by large black dots, known peculiar sources by small black dots, candidate binaries by red dots and known (resolved) binaries by encircled red dots (those listed in Table 3 are also labeled). Selection spaces, with inflection points listed in Table 5, are indicated by dashed lines




Fig. 5.— Best-fit composites for the resolved L dwarf/T dwarf binaries shown in Figure 3. In each panel, source spectra (black lines, in  $F_\lambda$  flux units) are shown normalized to the peak flux in the 1.0–1.3  $\mu\text{m}$  range; composites (green lines) are relatively normalized to minimize  $\chi^2$  deviations; primary (red lines) and secondary (blue lines) component spectra are normalized to their respective contribution to the composite (based on the faint Liu et al. 2006 relation; see Section 4.1). Noise spectra for candidate data are also indicated by grey lines. Component names, literature and index-based spectral types (the latter in parentheses) and  $\chi^2$  deviations are listed in the top right corner of each panel.




Fig. 6.— Spectral fits to the sources identified as “strong” binary candidates (at least three selection criteria satisfied). For each row, the left panel shows the best-fit single template (red lines) compared to the source spectrum (black lines, noise spectrum in grey), while the right panel shows the best-fit composite (green lines), primary (red lines) and secondary spectra (blue lines). Source spectra are normalized to the peak flux in the 1.0–1.3  $\mu\text{m}$  range, while single and composites are normalized to minimize  $\chi^2$  deviations. Component spectra in the right panels are normalized to their respective contribution to the composite, according to the faint Liu et al. (2006) relation (see Section 4.1). Template source names, literature and index-based spectral types (the latter in parentheses) and corresponding  $\chi^2$  deviations are labeled in the top right corner of each panel.




Fig. 6.— Continued.




Fig. 6.— Continued.






Fig. 6.— Continued.




Fig. 7.— Same as Figure 6 but for “weak” binary candidates (satisfying only 2 selection criteria).




Fig. 7.— Continued.




Fig. 7.— Continued.




Fig. 8.— Spectral fits for SDSS J1039+3256 (top), SDSS J1207+0244 (middle) and 2MASS J1324+6358 (bottom) based on flux calibrations with the faint (left) or bright (right) absolute magnitude relations of Liu et al. (2006). These were the only three sources for which the two relations gave statistically distinct  $\chi^2$  residuals. Format is the same as the right panels in Figures 6 and 7.




Fig. 9.— Absolute MKO  $J$  magnitudes versus spectral type for L and T dwarfs with parallax measurements. Spectral types are based on optical classifications for L2-L8 dwarfs (Kirkpatrick et al. 1999) and near-infrared classifications for L9-T8 dwarfs (Burgasser et al. 2006b). Known multiples (including the candidate 2MASS J1711+2232) and unresolved sources with absolute magnitude uncertainties  $\leq 0.3$  mag are indicated as open and filled circles, respectively. Absolute magnitudes for the primary (red) and secondary (blue) components of SDSS J0423-0414, SDSS J1021-0304, 2MASS J1711+2232 and  $\epsilon$  Indi BC (McCaughrean et al. 2004; Burgasser et al. 2006c; this paper) are also indicated by filled circles and connected to their combined-light magnitude and spectral types (dotted lines). The bright (top) and faint (bottom) absolute magnitude relations of Liu et al. (2006) are delineated by solid blue and red lines, respectively.

Table 1. New SpeX Spectral Observations.

Source Name (1)	Opt SpT (2)	NIR SpT (3)	2MASS $J$ (4)	UT Date (5)	Observer (6)	$t_{int}$ (s) (7)	Slit (") (8)	Airmass (9)	Cal Star (10)
2MASS J00165953-4056541	L3.5	...	15.32 $\pm$ 0.06	2008 Sep 7	AJB	900	0.5	2.04	HD 2339
2MASS J00193927-3724392	L3:	...	15.52 $\pm$ 0.06	2003 Sep 4	KLC	720	0.8	1.26	HD 9485
2MASS J0028208+224905	...	L7:	15.61 $\pm$ 0.07	2003 Sep 3	KLC	600	0.8	1.09	HD 198070
2MASSW J0030300-145033	L7	...	16.28 $\pm$ 0.11	2007 Jul 28	DLL	960	0.5	1.22	HD 1154
SDSSp J003259.36+141036.6	...	L8	16.83 $\pm$ 0.17	2008 Jul 13	AJB	900	0.5	1.02	HD 7215
2MASSW J0051107-154417	L3.5	...	15.28 $\pm$ 0.05	2008 Sep 8	AJB	960	0.5	1.24	HD 1154
2MASS J00531899-3631102	L3.5	...	14.45 $\pm$ 0.03	2008 Sep 7	AJB	360	0.5	1.83	HD 5061
SDSSp J010752.33+004156.1	L8	L5.5	15.82 $\pm$ 0.06	2007 Oct 12	DLL	1200	0.5	1.16	HD 9485
2MASS J01311838+3801554	L4:	...	14.68 $\pm$ 0.03	2003 Sep 3	KLC	720	0.8	1.27	HD 7215
2MASS J01550354+0950003	...	L5:	14.83 $\pm$ 0.04	2003 Sep 4	KLC	360	0.8	1.05	BD+18 337A
DENIS-P J0205.4-1159	L7	L5.5 $\pm$ 2	14.59 $\pm$ 0.03	2005 Sep 7	AJB	720	0.5	1.42	HD 13936
2MASSW J0208236+273740	L5	...	15.71 $\pm$ 0.06	2008 Sep 7	AJB	600	0.5	1.11	HD 19600
2MASS J03185403-3421292	L7	...	15.57 $\pm$ 0.06	2003 Sep 3	KLC	1440	0.8	1.70	HD 22861
2MASS J0447430-193604	L5:	...	15.97 $\pm$ 0.07	2003 Sep 4	KLC	1080	0.8	1.52	HD 20423
2MASS J05161597-3332046	L3:	...	15.88 $\pm$ 0.06	2004 Nov 7	KLC	720	0.5	1.75	HD 31506
2MASSI J0652307+471034	L4.5	...	13.51 $\pm$ 0.02	2004 Nov 7	KLC	720	0.5	1.17	HD 45105
SDSS J065405.63+652805.4	...	L6 $\pm$ 1	16.14 $\pm$ 0.09	2008 Jan 8	AJB	1440	0.5	1.46	HD 38831
2MASSW J0717163+570543	...	L6.5	14.64 $\pm$ 0.03	2008 Jan 8	AJB	480	0.5	1.30	HD 38831
SDSS J073922.26+661503.5	...	T1.5 $\pm$ 1	16.82 $\pm$ 0.13	2008 Jan 9	AJB	900	0.5	1.47	HD 48049
SDSS J074149.15+235127.5	...	T5	16.16 $\pm$ 0.10	2008 Jan 10	AJB	1800	0.5	1.00	HD 58296
SDSS J074201.41+205520.5	...	T5	16.19 $\pm$ 0.09	2008 Jan 10	AJB	1800	0.5	1.02	HD 58296
2MASSW J0801405+462850	L6.5	...	16.28 $\pm$ 0.13	2008 Jan 12	AJB	1500	0.5	1.13	HD 63160
2MASSW J0820299+450031	L5	...	16.28 $\pm$ 0.11	2007 Nov 13	KLC	900	0.5	1.11	HD 71906
2MASS J08234818+2428577	L3	...	14.99 $\pm$ 0.04	2008 Jan 10	AJB	480	0.5	1.03	HD 74721
2MASSI J0825196+211552	L7.5	L6	15.10 $\pm$ 0.03	2005 Mar 23	KLC	1080	0.5	1.01	HD 58729
SDSS J083048.80+012831.1	...	T4.5	16.29 $\pm$ 0.11	2008 Jan 10	AJB	900	0.5	1.11	HD 74721
2MASSI J0835425-081923	L5	...	13.17 $\pm$ 0.02	2004 Nov 8	KLC	360	0.5	1.24	HD 79752
SDSS J085234.90+472035.0	...	L9.5 $\pm$ 1	16.18 $\pm$ 0.11	2008 Jan 9	AJB	1200	0.5	1.13	HD 63160
2MASSI J0856479+223518	L3:	...	15.68 $\pm$ 0.07	2007 Nov 13	KLC	1080	0.5	1.04	HD 74721
SDSSp J085758.45+570851.4	L8	L8 $\pm$ 1	15.04 $\pm$ 0.04	2008 Jan 8	AJB	960	0.5	1.25	HD 38831
SDSS J085834.42+325627.7	...	T1	16.45 $\pm$ 0.12	2008 Jan 9	AJB	1200	0.5	1.05	HD 71906
2MASS J09054654+5623117	...	L5	15.40 $\pm$ 0.05	2004 Nov 7	KLC	720	0.5	1.36	HD 71906
Gliese 337CD	L8	T0	15.51 $\pm$ 0.08	2008 Jan 12	AJB	900	0.5	1.03	HD 74721

Table 1—Continued

Source Name (1)	Opt SpT (2)	NIR SpT (3)	2MASS $J$ (4)	UT Date (5)	Observer (6)	$t_{int}$ (s) (7)	Slit (") (8)	Airmass (9)	Cal Star (10)
2MASSW J0920122+351742	L6.5	T0p	15.63±0.06	2008 Jan 9	AJB	1200	0.5	1.08	HD 71906
SDSSp J092615.38+584720.9	...	T4.5	15.90±0.07	2007 Nov 11	DLL	1800	0.5	1.46	HD 33654
2MASSW J0929336+342952	L8	L7.5	16.60±0.13	2008 Jan 9	AJB	150	0.5	1.10	HD 89239
SDSS J100711.74+193056.2	...	L8±1.5	16.87±0.16	2008 Jan 12	AJB	1500	0.5	1.05	HD 89239
SDSS J103026.78+021306.4	...	L9.5±1	>16.89	2008 Jan 10	AJB	1800	0.5	1.10	HD 96781
2MASSW J1036530-344138	L6	...	15.62±0.05	2008 Jan 13	AJB	900	0.5	1.86	HD 86382
SDSS J103931.35+325625.5	...	T1	16.41±0.15	2008 Jan 13	AJB	1500	0.5	1.22	HD 89239
SDSS J104335.08+121314.1	...	L7±1	16.00±0.09	2008 Jan 12	AJB	720	0.5	1.09	HD 101060
SDSS J104409.43+042937.6	...	L7	15.88±0.08	2008 Jan 10	AJB	900	0.5	1.17	HD 96781
SDSS J104829.21+091937.8	...	T2.5	16.59±0.15	2008 Jan 13	AJB	1800	0.5	1.07	HD 101060
DENIS-P J1058.7-1548	L3	L3	14.16±0.04	2005 Mar 24	KLC	720	0.5	1.13	HD 93346
Gliese 417BC	L4.5	...	14.58±0.03	2006 Dec 23	AJB	720	0.5	1.04	HD 89239
2MASSW J1146345+223053	L3	...	14.17±0.03	2008 Jan 12	AJB	360	0.5	1.04	HD 101060
SDSS J115553.86+055957.5	...	L7.5	15.66±0.08	2008 Jan 13	AJB	900	0.5	1.11	HD 101060
2MASS J12070374-3151298	...	L3:	15.85±0.07	2008 Jan 9	AJB	960	0.5	1.62	HD 105379
SDSS J121951.45+312849.4	...	L8	15.91±0.08	2006 Dec 24	AJB	900	0.5	1.08	HD 99966
DENIS-P J1228.2-1547	L5	L6±2	14.38±0.03	2008 Jan 14	AJB	1200	0.5	1.28	HD 105764
2MASSW J1239272+551537	L5	...	14.71±0.03	2008 Jan 11	AJB	480	0.5	1.23	HD 99966
SDSSp J132629.82-003831.5	L8:	L5.5	16.10±0.07	2005 Mar 24	KLC	1080	0.5	1.28	HD 111744
SDSS J133148.92-011651.4	L6	L8±2.5	15.46±0.04	2008 Jan 14	AJB	1500	0.5	1.12	HD 123233
2MASS J13571237+1428398	L4:	...	15.58±0.06	2005 Mar 24	KLC	1080	0.5	1.05	HD 121880
SDSS J140023.12+433822.3	...	L7±1	16.30±0.09	2008 Jul 13	AJB	1200	0.5	1.13	HD 128039
2MASS J14075361+1241099	L5	...	15.38±0.06	2008 Jul 30	AJB	900	0.5	1.17	HD 122945
SDSS J141530.05+572428.7	...	T3±1	16.73±0.16	2008 Jul 14	AJB	1500	0.5	1.46	HD 116405
2MASS J14283132+5923354	...	L4	14.78±0.04	2003 Sep 5	KLC	360	0.8	1.70	HD 145454
2MASSW J1448256+103159	...	L3.5	14.56±0.03	2005 Mar 23	KLC	900	0.5	1.03	HD 124773
SDSS J151114.66+060742.9	...	T0±2	16.02±0.08	2007 Jul 28	DLL	960	0.5	1.15	HD 136831
2MASSW J1515008+484742	L6	L6	14.11±0.03	2003 Sep 5	KLC	360	0.8	1.45	HD 145454
SDSS J151506.11+443648.3	...	L7.5±1.5	16.58±0.15	2008 Jul 14	AJB	900	0.5	1.24	HD 143187
SDSS J151603.03+025928.9	...	T0:	17.23±0.20	2008 Jul 13	AJB	1440	0.5	1.19	HD 136831
SDSS J151643.01+305344.4	...	T0.5±1	16.85±0.15	2008 Jul 12	AJB	1050	0.5	1.24	HD 127304
SDSS J152039.82+354619.8	...	T0±1	15.54±0.06	2008 Jul 12	AJB	900	0.5	1.15	HD 127304
SDSS J152103.24+013142.7	...	T2:	16.40±0.10	2008 Jul 13	AJB	1440	0.5	1.36	HD 136831



Table 1—Continued

Source Name (1)	Opt SpT (2)	NIR SpT (3)	2MASS $J$ (4)	UT Date (5)	Observer (6)	$t_{int}$ (s) (7)	Slit (") (8)	Airmass (9)	Cal Star (10)
Gliese 584C	L8	L8	16.06±0.10	2008 May 8	AJB	900	0.5	1.05	HD 140729
2MASS J1534498-295227	T6	T5.5	14.90±0.05	2006 Mar 12	AJB	1800	0.8	1.55	HD 138295
DENIS-P J153941.96-052042.4	L4:	L2	13.92±0.03	2008 Jul 14	AJB	360	0.5	1.11	HD 137873
SDSS J154009.36+374230.3	...	L9±1.5	16.56±0.14	2008 Jul 30	AJB	1440	0.5	1.40	HD 140729
2MASS J15461461+4932114	...	T2.5±1	15.90±0.07	2008 Jul 12	AJB	600	0.5	1.50	HD 143187
2MASS J1553022+153236	...	T7	15.83±0.07	2006 Apr 8	AJB	1440	0.5	1.17	HD 140729
SDSS J163030.53+434404.0	...	L7±1.5	16.63±0.14	2008 Jul 13	AJB	1200	0.5	1.38	HD 1533415
SDSS J163239.34+415004.3	...	T1:	17.08±0.22	2008 Jul 14	AJB	1200	0.5	1.14	HD 143187
2MASS J16452207+3004071	L3	...	15.19±0.04	2008 Jul 13	AJB	480	0.5	1.70	HD 1533415
SDSS J165329.69+623136.5	L3	...	15.09±0.05	2008 Jul 30	AJB	480	0.5	1.67	HD 143187
DENIS-P J170548.38-051645.7	...	L4	13.31±0.03	2004 Sep 7	AJB	540	0.5	1.30	HD 159415
2MASS J1711457+223204	L6.5	...	17.09±0.18	2008 Sep 7	AJB	1500	0.5	1.05	HD 165029
2MASS J17461199+5034036	...	L5	15.10±0.06	2003 Sep 4	KLC	720	0.8	1.26	HD 172728
2MASS J17502484-0016151	...	L5.5	13.29±0.02	2007 Sep 16	AJB	540	0.5	1.07	HD 161289
2MASS J19285196-4356256	L5	...	15.20±0.04	2008 Sep 8	AJB	1320	0.5	2.28	HD 186216
2MASS J20343769+0827009	L3	...	14.46±0.03	2008 Jul 29	AJB	600	0.5	1.05	HD 189920
2MASS J20360316+1051295	L3	...	13.95±0.03	2008 Jul 29	AJB	540	0.5	1.02	HD 189920
SDSS J204317.69-155103.4	...	L9	16.63±0.16	2008 Jul 12	AJB	1440	0.5	1.23	HD 200604
SDSS J204749.61-071818.3	...	T0:	16.95±0.20	2008 Jul 13	AJB	1200	0.5	1.16	HD 203769
SDSS J205235.31-160929.8	...	T1±1	16.33±0.12	2008 Jul 12	AJB	1200	0.5	1.27	HD 200604
2MASSW J2101154+175658	L7.5	...	16.85±0.17	2008 Jul 12	AJB	1200	0.5	1.06	GSC 02716-02882
2MASS J2104149-103736	L3	...	13.84±0.03	2003 Sep 5	KLC	240	0.8	1.05	HD 198070
2MASS J21373742+0808463	L4:	...	14.77±0.03	2007 Nov 14	KLC	1080	0.5	1.37	HD 208108
2MASS J21395208+2148397	L3:	...	15.88±0.07	2003 Sep 3	KLC	720	0.8	1.02	HD 198070
SDSS J214046.55+011259.7	L3	...	15.89±0.08	2008 Sep 8	AJB	900	0.5	1.14	HD 196442
2MASS J21420580-3101162	L3	...	15.84±0.07	2008 Sep 8	AJB	1440	0.5	1.69	HD 200835
2MASS J22120703+3430351	...	L5:	16.32±0.10	2003 Sep 4	KLC	2160	0.8	1.10	HD 210290
2MASSW J2224438-015852	L4.5	L3.5	14.07±0.03	2003 Sep 5	KLC	360	0.8	1.10	HD 215143
2MASS J22425317+2542573	L3	...	14.81±0.04	2008 Jul 12	AJB	720	0.5	1.05	GSC 01149-00254
2MASS J23254530+4251488	L8	...	15.49±0.05	2003 Aug 12	KLC	720	0.8	1.09	HD 6574
2MASS J2325560-025950	L3:	...	15.96±0.08	2003 Sep 3	KLC	1440	0.8	1.26	HD 213030
2MASS J23392527+3507165	...	L3.5	15.36±0.05	2003 Sep 4	KLC	720	0.8	1.16	HD 210290



Table 2. Spectral Templates.

Source Name (1)	Opt SpT (2)	NIR SpT (3)	SpeX SpT (4)	2MASS $J$ (5)	$J - K_s$ (6)	Ref <sup>a</sup> (7)
SDSS J000013.54+255418.6	...	T4.5	T4.5	15.06±0.04	0.23±0.13	1; 2,1
2MASS J0013578-223520	L4	...	L3.5	15.78±0.07	1.74±0.08	3; 4
2MASS J00165953-4056541	L3.5	...	L4.5±1	15.32±0.06	1.88±0.07	3; 5
2MASS J00193927-3724392	L3:	...	L3.5±1	15.52±0.06	1.83±0.08	3; 6
2MASS J0028208+224905 <sup>d</sup>	...	L7:	L5.0	15.61±0.07	1.83±0.10	3; 6
2MASSW J0030300-145033 <sup>d</sup>	L7	...	L4.5±2	16.28±0.11	1.80±0.15	3; 7
SDSSp J003259.36+141036.6	...	L8	L7.5	16.83±0.17	1.88±0.20	3; 8
2MASS J00345157+0523050	...	T6.5	T6.5	15.54±0.05	<-0.71	9; 9,1
2MASSW J0036159+182110	L3.5	L4±1	L4.0	12.47±0.03	1.41±0.03	10; 11,7,2
HD 3651B	...	T7.5	T8.0	16.16±0.03	-0.71±0.06	12; 13,14
2MASS J00501994-3322402	...	T7	T6.5	15.93±0.07	0.69±0.20	15; 16,1
2MASSW J0051107-154417	L3.5	...	L5.0±1	15.28±0.05	1.81±0.06	3; 7
2MASS J00531899-3631102	L3.5	...	L4.0	14.45±0.03	1.51±0.04	3; 5
2MASS J0103320+193536	L6	...	L6.0	16.29±0.08	2.14±0.10	17; 7
SDSSp J010752.33+004156.1 <sup>d</sup>	L8	L5.5	L6.0	15.82±0.06	2.12±0.07	3; 8,18,2
SDSS J011912.22+240331.6 <sup>c</sup>	...	T2	T2.5	17.02±0.18	<-0.02	23; 23
2MASS J01311838+3801554 <sup>d</sup>	L4:	...	L1.5	14.68±0.03	1.63±0.05	3; 19
IPMS J013656.57+093347.3	...	T2.5	T2.0	13.46±0.03	0.89±0.04	10; 20
2MASS J01443536-0716142	L5	...	L4.5	14.19±0.03	1.92±0.03	10; 21
SDSS J015141.69+124429.6 <sup>b</sup>	...	T1	T0.0±1	16.57±0.13	1.38±0.23	9; 8,1
2MASS J01550354+0950003	...	L5:	L5.0	14.83±0.04	1.69±0.05	3; 6
2MASSW J0205034+125142	L5	...	L5.5	15.68±0.06	2.01±0.07	22; 7
SDSS J020608.97+223559.2	...	L5.5	L6.0	16.56±0.11	1.39±0.17	23; 23
2MASSW J0208236+273740	L5	...	L5.0	15.71±0.06	1.84±0.08	3; 7
2MASS J0243137-245329	...	T6	T6.0	15.38±0.05	0.16±0.18	9; 24,1
SDSS J024749.90-163112.6 <sup>c</sup>	...	T2±1.5	T2.5±1	17.19±0.18	1.57±0.27	23; 23
DENIS-P J0255-4700	L8	L9	L9.0±2	13.25±0.03	1.69±0.04	1; 25,5,1
2MASS J02572581-3105523	L8	...	L6.0	14.67±0.04	1.80±0.05	26; 5
2MASS J03185403-3421292	L7	...	L6.5	15.57±0.06	2.06±0.07	3; 5
SDSS J032553.17+042540.1	...	T5.5	T5.5	16.25±0.14	<-0.27	23; 23
2MASS J0328426+230205	L8	L9.5	L9.0±1	16.69±0.14	1.78±0.18	10; 7,2
SDSS J035104.37+481046.8 <sup>c</sup>	...	T1±1.5	T1.5±1	16.47±0.13	1.47±0.18	23; 23
2MASS J04070885+1514565	...	T5	T5.5	16.06±0.09	0.13±0.28	9; 9,9
2MASS J0415195-093506	T8	T8	T8.0	15.70±0.06	0.27±0.21	9; 24,27,1
2MASS J0439010-235308	L6.5	...	L6.0	14.41±0.03	1.59±0.04	28; 29
2MASS J0447430-193604	L5:	...	L4.0±1	15.97±0.07	1.96±0.08	3; 6
2MASS J05103520-4208140	...	T5	T5.5	16.22±0.09	0.23±0.30	30; 30
2MASS J05160945-0445499	...	T5.5	T5.5	15.98±0.08	0.50±0.22	10; 31,1
2MASS J05161597-3332046	L3:	...	L4.5±1	15.88±0.06	1.89±0.07	3; 6
2MASS J05591914-1404488	T5	T4.5	T4.5	13.80±0.02	0.23±0.06	1; 327,1
2MASS J06020638+4043588	...	T4.5	T4.5	15.54±0.07	0.38±0.18	30; 30
2MASS J06244595-4521548	L5:	...	L5.0	14.48±0.03	1.88±0.04	26; 33
2MASS J0652307+471034	L4.5	...	L6.5	13.51±0.02	1.82±0.03	3; 29
SDSS J065405.63+652805.4	...	L6±1	L5.0	16.14±0.09	1.54±0.12	3; 23

Table 2—Continued

Source Name (1)	Opt SpT (2)	NIR SpT (3)	SpeX SpT (4)	2MASS $J$ (5)	$J - K_s$ (6)	Ref <sup>a</sup> (7)
2MASSW J0717163+570543	...	L6.5	L5.0	14.64±0.03	1.69±0.04	3; 34
2MASSI J0727182+171001	T8	T7	T7.0	15.60±0.06	0.04±0.20	15; 24,27,1
SDSS J073922.26+661503.5	...	T1.5±1	T2.0	16.82±0.13	0.99±0.22	3; 23
SDSS J074149.15+235127.5	...	T5	T5.5	16.16±0.10	<0.31	3; 2,1
SDSS J074201.41+205520.5	...	T5	T5.0	16.19±0.09	<0.97	3; 2,1
2MASSI J0755480+221218	T6	T5	T5.0	15.73±0.06	-0.03±0.22	1; 24,27,1
SDSS J075840.33+324723.4 <sup>c</sup>	...	T2	T2.5	14.95±0.04	1.07±0.07	10; 2,1
2MASSW J0801405+462850	L6.5	...	L5.5	16.28±0.13	1.74±0.17	3; 7
2MASSW J0820299+450031	L5	...	L4.5±2	16.28±0.11	2.06±0.13	3; 7
2MASS J08234818+2428577	L3	...	L1.5	14.99±0.04	1.61±0.05	3; 33
2MASSI J0825196+211552	L7.5	L6	L7.0	15.10±0.03	2.07±0.04	3; 7,2
SDSS J083048.80+012831.1	...	T4.5	T5.0	16.29±0.11	<-0.07	3; 2,1
SDSSp J083008.12+482847.4	L8	L9±1	L9.5±1	15.44±0.05	1.77±0.06	10; 8,5
SDSS J083506.16+195304.4	...	L4.5	L6.0	16.09±0.08	1.77±0.09	23; 23
2MASSI J0835425-081923	L5	...	L5.0	13.17±0.02	2.03±0.03	3; 29
SDSS J085116.20+181730.0	...	L4.5±1.5	L5.5±1	16.57±0.12	1.60±0.16	23; 23
SDSS J085234.90+472035.0	...	L9.5±1	L9.0±1	16.18±0.11	1.46±0.16	3; 2
SDSSp J085758.45+570851.4 <sup>b</sup>	L8	L8±1	L8.5	15.04±0.04	2.08±0.05	3; 8,5
SDSS J085834.42+325627.7 <sup>b</sup>	...	T1	T0.0±2	16.45±0.12	1.70±0.15	23; 23
2MASS J09054654+5623117	...	L5	L6.0	15.40±0.05	1.67±0.06	3; 33
2MASSI J0908380+503208	L5	L9±1	L6.0	14.55±0.02	1.60±0.04	26; 29,2
SDSS J090900.73+652527.2 <sup>c</sup>	...	T1.5	T1.5	16.03±0.09	0.86±0.17	23; 23
2MASSW J0929336+342952	L8	L7.5	L7.5	16.60±0.13	1.96±0.16	3; 7,2
2MASSI J0937347+293142	T7	T6p	T5.5	14.65±0.04	-0.62±0.13	15; 24,27,1
2MASS J09490860-1545485 <sup>c</sup>	...	T2	T1.5±1	16.15±0.12	0.92±0.20	1; 16,1
SDSS J100711.74+193056.2	...	L8±1.5	L9.5	16.87±0.16	1.87±0.19	3; 23
2MASS J10073369-4555147	...	T5	T5.5	15.65±0.07	0.09±0.24	30; 30
2MASSI J1010148-040649	L6	...	L5.0	15.51±0.06	1.89±0.07	22; 29
SDSS J103026.78+021306.4	...	L9.5±1	L9.0±1	17.10±0.05 <sup>e</sup>	1.43±0.07 <sup>e</sup>	3; 2
2MASSW J1036530-344138	L6	...	L6.5	15.62±0.05	1.82±0.06	3; 35
SDSS J103931.35+325625.5 <sup>c</sup>	...	T1	T1.5	16.41±0.15	1.25±0.22	3; 23
SDSS J104335.08+121314.1	...	L7±1	L8.0	16.00±0.09	1.74±0.11	3; 23
2MASS J10430758+2225236	L8	...	L9.0±1	15.97±0.07	1.97±0.08	26; 19
SDSS J104409.43+042937.6	...	L7	L7.0	15.88±0.08	1.62±0.12	3; 2
2MASSI J1047538+212423	T7	T6.5	T6.0	15.82±0.06	<-0.59	10; 36,27,1
SDSS J104829.21+091937.8	...	T2.5	T2.5	16.59±0.15	<0.23	3; 23
DENIS-P J1058.7-1548	L3	L3	L3.0	14.16±0.04	1.62±0.05	3; 37,38,2
2MASS J10595185+304205	...	T4	T4.0	16.20±0.09	0.64±0.20	39; 39
2MASS J11000965+4957470	...	L3.5	L3.5	15.28±0.04	1.81±0.05	26; 33
2MASSI J1104012+195921	L4	...	L5.5	14.38±0.03	1.43±0.04	9; 29
2MASS J11061197+2754225 <sup>c</sup>	...	T2.5	T2.0	14.82±0.04	1.02±0.07	30; 30
2MASS J11145133-2618235	...	T7.5	T7.5	15.86±0.08	<-0.25	15; 16,1
2MASS J11220826-3512363 <sup>b</sup>	...	T2	T2.0	15.02±0.04	0.64±0.08	1; 16,1
SDSS J115553.86+055957.5	...	L7.5	L7.0	15.66±0.08	1.54±0.11	3; 2

Table 2—Continued

Source Name (1)	Opt SpT (2)	NIR SpT (3)	SpeX SpT (4)	2MASS $J$ (5)	$J - K_s$ (6)	Ref <sup>a</sup> (7)
SDSS J120602.51+281328.7 <sup>c</sup>	...	T3	T3.0	16.54±0.11	<0.72	23; 23
SDSS J120747.17+024424.8 <sup>c</sup>	L8	T0	T0.0	15.58±0.07	1.59±0.09	30; 18,1
2MASS J12070374-3151298	...	L3:	L2.5	15.85±0.07	1.85±0.09	26; 33
2MASS J12095613-1004008 <sup>b</sup>	...	T3	T3.0	15.91±0.07	0.85±0.16	9; 9,1
SDSS J121440.95+631643.4 <sup>b</sup>	...	T3.5±1	T4.0	16.59±0.12	0.71±0.26	23; 23
2MASSI J1217110-031113	T7	T7.5	T7.5	15.86±0.06	<-0.03	15; 36,27,1
SDSS J121951.45+312849.4	...	L8	L9.5	15.91±0.08	1.61±0.11	3; 23
2MASS J12314753+0847331	...	T5.5	T6.0	15.57±0.07	0.35±0.21	9; 9,1
2MASS J12373919+6526148	T7	T6.5	T6.5	16.05±0.09	<-0.01	54; 36,27,1
SDSSp J125453.90-012247.4 <sup>b</sup>	T2	T2	T2.0	14.89±0.04	1.05±0.06	9; 40,27,1
SDSSp J132629.82-003831.5	L8:	L5.5	L6.0±2	16.10±0.07	1.90±0.10	3; 41,2
SDSSp J134646.45-003150.4	T7	T6.5	T6.0	16.00±0.10	0.23±0.29	15; 42,27,1
2MASS J13571237+1428398	L4:	...	L3.0±1	15.58±0.06	1.70±0.07	3; 6
SDSS J135852.68+374711.9	...	T4.5±1	T5.0	16.46±0.09	<0.36	23; 23
SDSS J140023.12+433822.3 <sup>d</sup>	...	L7±1	L5.0	16.30±0.09	1.81±0.12	3; 23
2MASS J14075361+1241099	L5	...	L5.0	15.38±0.06	1.78±0.07	3; 33
SDSS J141530.05+572428.7 <sup>c</sup>	...	T3±1	T3.0±1	16.73±0.16	<1.19	3; 23
2MASS J14283132+5923354	...	L4	L4.5	14.78±0.04	1.52±0.05	3; 33
SDSS J143553.25+112948.6 <sup>c</sup>	...	T2±1	T2.5±1	17.14±0.23	<0.23	23; 23
SDSS J143945.86+304220.6 <sup>c</sup>	...	T2.5	T2.5	17.22±0.23	<1.34	23; 23
2MASSW J1448256+103159 <sup>d</sup>	L4:	L3.5	L7.0	14.56±0.03	1.87±0.05	3; 33,34
Gliese 570D	T7	T7.5	T7.5	15.32±0.05	0.08±0.16	9; 43,27,1
2MASS J15031961+2525196	T6	T5	T5.5	13.94±0.02	-0.03±0.06	9; 44,27,1
2MASSW J1506544+132106	L3	...	L4.0	13.37±0.02	1.62±0.03	28; 45,45
2MASSW J1507476-162738	L5	L5.5	L5.0	12.83±0.03	1.52±0.04	28; 7,2
SDSS J151506.11+443648.3	...	L7.5±1.5	L7.0	16.58±0.15	1.83±0.20	3; 23
2MASSW J1515008+484742	L6	L6	L5.5	14.11±0.03	1.61±0.04	3; 34,19
SDSS J151603.03+025928.9 <sup>c</sup>	...	T0:	L9.5±1	17.23±0.20	1.80±0.27	3; 2,1
SDSS J152039.82+354619.8 <sup>d</sup>	...	T0±1	L7.5	15.54±0.06	1.54±0.08	3; 23
SDSS J152103.24+013142.7	...	T2:	T2.5	16.40±0.10	1.05±0.20	3; 2,1
Gliese 584C	L8	L8	L7.5	16.06±0.10	1.71±0.12	3; 7,8
2MASSI J1526140+204341	L7	...	L5.5	15.59±0.06	1.66±0.08	9; 7
DENIS-P J153941.96-052042.4	L4:	L2	L4.0	13.92±0.03	1.35±0.04	3; 46,5
SDSS J154009.36+374230.3	...	L9±1.5	L9.0±1	16.56±0.14	1.82±0.16	3; 23
2MASS J15461461+4932114	...	T2.5±1	T3.0	15.90±0.07	<1.00	3; 47
2MASSI J1546291-332511	...	T5.5	T5.5	15.63±0.05	0.15±0.19	10; 24,1
2MASS J16150413+1340079	...	T6	T6.0	16.35±0.09	<0.49	30; 30
SDSSp J162414.37+002915.6	...	T6	T6.0	15.49±0.05	<-0.02	15; 48,1
SDSS J163030.53+434404.0	...	L7±1.5	L7.5	16.63±0.14	1.98±0.16	3; 2
2MASSW J1632291+190441	L8	L8	L7.0	15.87±0.07	1.86±0.08	28; 38,1
SDSS J163239.34+415004.3 <sup>b</sup>	...	T1:	T1.0	17.08±0.22	<1.33	3; 2,1
SDSS J163359.23-064056.5	...	L6	L5.5	16.14±0.09	1.59±0.13	23; 23
2MASS J16452207+3004071	L3	...	L1.5	15.19±0.04	1.60±0.05	3; 19
SDSS J165329.69+623136.5	L3	...	L1.0	15.09±0.05	<1.02	3; 18

Table 2—Continued

Source Name (1)	Opt SpT (2)	NIR SpT (3)	SpeX SpT (4)	2MASS $J$ (5)	$J - K_s$ (6)	Ref <sup>a</sup> (7)
DENIS-P J170548.38-051645.7 <sup>d</sup>	L0.5	L4	L1.0	13.31±0.03	1.28±0.04	3; 33,46
2MASS J1711457+223204 <sup>c,d</sup>	L6.5	...	L9.0±3	17.09±0.18	2.36±0.20	3; 7
2MASS J17461199+5034036	...	L5	L5.0	15.10±0.06	1.57±0.07	3; 33
SDSSp J175032.96+175903.9	...	T3.5	T3.0	16.34±0.10	0.86±0.21	9; 8,1
SDSS J175024.01+422237.8 <sup>b</sup>	...	T2	T1.5	16.47±0.10	0.98±0.20	1; 2,1
2MASS J17502484-0016151	...	L5.5	L6.5	13.29±0.02	1.44±0.03	3; 49
2MASS J1754544+164920	...	T5	T5.5	15.76±0.09	<0.97	10; 50
SDSS J175805.46+463311.9	...	T6.5	T6.5	16.15±0.09	0.69±0.21	15; 2,1
2MASS J18283572-4849046	...	T5.5	T5.5	15.18±0.06	-0.01±0.15	9; 9,1
2MASS J19010601+4718136	...	T5	T5.0	15.86±0.07	0.22±0.29	9; 9,1
2MASS J19285196-4356256	L5	...	L5.5	15.20±0.04	1.74±0.06	3; 33
SDSS J202820.32+005226.5	L3	...	L2.0	14.30±0.04	1.51±0.05	9; 18
2MASS J20343769+0827009 <sup>d</sup>	L3	...	L0.5	14.46±0.03	1.38±0.05	3; 33
2MASS J20360316+1051295	L3	...	L3.5	13.95±0.03	1.50±0.04	3; 33
SDSS J204317.69-155103.4	...	L9	L9.0±1	16.63±0.16	1.22±0.27	3; 23
SDSS J204749.61-071818.3 <sup>d</sup>	...	T0:	L7.5	16.95±0.20	<1.96	3; 2,1
SDSS J205235.31-160929.8 <sup>c</sup>	...	T1±1	T0.0±1	16.33±0.12	1.21±0.19	3; 23
2MASS J2104149-103736	L3	...	L2.0	13.84±0.03	1.47±0.04	3; 29
SDSS J212413.89+010000.3	...	T5	T5.0	16.03±0.07	<-0.11	1; 2,1
SDSS J213154.43-011939.3	...	L9	L8.5±1	17.40±0.26	1.84±0.34	23; 23
2MASS J21321145+1341584	L6	...	L5.0	15.80±0.06	1.96±0.08	26; 19
2MASS J21373742+0808463	L5:	...	L5.0	14.77±0.03	1.76±0.04	3; 33
2MASS J21392676+0220226 <sup>c</sup>	T0	T1.5	T2.5±1	15.26±0.05	1.68±0.07	1; 33,1
2MASS J21395208+2148397	L3:	...	L2.0	15.88±0.07	1.82±0.10	3; 6
SDSS J214046.55+011259.7	L3	...	L4.5	15.89±0.08	1.47±0.11	3; 18
2MASS J21420580-3101162	L3	...	L2.0	15.84±0.07	1.88±0.08	3; 51
HN Peg B <sup>b</sup>	...	T2.5	T3.0	15.86±0.03	0.74±0.04	14; 14
2MASS J21512543-2441000 <sup>d</sup>	L3	...	L5.0	15.75±0.08	2.10±0.09	10; 51,19
2MASS J21513839-4853542	...	T4	T4.0	15.73±0.07	0.30±0.20	1; 52,1
2MASS J21543318+5942187	...	T5	T5.5	15.66±0.07	<0.32	30; 30
2MASS J21542494-1023022	...	T4.5	T4.5	16.42±0.12	<-0.62	30; 30
2MASS J21580457-1550098	L4:	...	L5.0	15.04±0.04	1.85±0.05	55; 5
2MASS J22120703+3430351	...	L5:	L5.5	16.32±0.10	1.95±0.12	3; 33
2MASSW J2224438-015852	L4.5	L3.5	L4.5±1	14.07±0.03	2.05±0.04	3; 7,2
2MASS J22282889-4310262	...	T6	T6.0	15.66±0.07	0.37±0.22	9; 31,1
2MASS J22425317+2542573	L3	...	L1.5	14.81±0.04	1.76±0.05	3; 53,19
2MASS J2254188+312349	...	T4	T4.0	15.26±0.05	0.36±0.15	9; 24,1
2MASS J23254530+4251488	L8	...	L7.5±1	15.49±0.05	1.73±0.07	3; 19
2MASS J2325560-025950	L3:	...	L3.0±1	15.96±0.08	1.85±0.10	3; 6
2MASS J23312378-4718274	...	T5	T5.0	15.66±0.07	0.27±0.21	9; 9,1
2MASS J2339101+135230	...	T5	T5.0	16.24±0.11	0.09±0.33	1; 24,1
2MASS J23392527+3507165	...	L3.5	L4.5	15.36±0.05	1.77±0.07	3; 33
2MASS J2356547-155310	...	T5.5	T5.5	15.82±0.06	0.05±0.19	1; 24,1

<sup>a</sup>Data reference followed by references for discovery, optical spectral type and near-infrared spectral type, where available.

<sup>b</sup>Source satisfied one spectral index selection criterion, used as a template for spectral fitting.

<sup>c</sup>Source satisfied two or more spectral index selection criteria (Table 7) and not used as a template for spectral fitting.

<sup>d</sup>Optical, near-infrared and/or SpeX classifications deviate by two subtypes or more.

<sup>e</sup>Undetected in 2MASS; photometry on the Mauna Kea Observatory (MKO) system (Knapp et al. 2004).

References. — (1) Burgasser et al. (2006b); (2) Knapp et al. (2004); (3) This paper; (4) Kendall et al. (2003); (5) Kirkpatrick et al. (2008); (6) K. Cruz, in preparation; (7) Kirkpatrick et al. (2000); (8) Geballe et al. (2002); (9) Burgasser et al. (2004); (10) Burgasser et al. (2008a); (11) Reid et al. (2000); (12) Burgasser (2007c); (13) Mugrauer et al. (2006); (14) Luhman et al. (2007); (15) Burgasser et al. (2006a); (16) Tinney et al. (2005); (17) Cruz et al. (2004); (18) Schneider et al. (2002); (19) Cruz et al. (2007); (20) Artigau et al. (2006); (21) Liebert et al. (2003); (22) Reid et al. (2006); (23) Chiu et al. (2006); (24) Burgasser et al. (2002a); (25) Martín et al. (1999); (26) Siegler et al. (2007); (27) Burgasser et al. (2003b); (28) Burgasser (2007a); (29) Cruz et al. (2003); (30) Looper et al. (2007); (31) Burgasser et al. (2003d); (32) Burgasser et al. (2000a); (33) Reid et al. (2008); (34) Wilson et al. (2003); (35) Gizis (2002); (36) Burgasser et al. (1999); (37) Delfosse et al. (1997); (38) Kirkpatrick et al. (1999); (39) Sheppard & Cushing (2009); (40) Leggett et al. (2000); (41) Fan et al. (2000); (42) Tsvetanov et al. (2000); (43) Burgasser et al. (2000b); (44) Burgasser et al. (2003c); (45) Gizis et al. (2000); (46) Kendall et al. (2004); (47) Metchev et al. (2008); (48) Strauss et al. (1999); (49) Kendall et al. (2007); (50) A. Burgasser, in preparation; (51) Liebert & Gizis (2006); (52) Ellis et al. (2005); (53) Gizis et al. (2003); (54) Liebert & Burgasser (2007); (55) J. D. Kirkpatrick, in preparation.

Table 3. Reference L Dwarf/T Dwarf Transition Binaries.

Source Name (1)	Opt SpT (2)	NIR SpT (3)	Component SpTs (4)	Component Relative Magnitudes <sup>a</sup> (5)		Ref (6)
SDSSp J042348.57-041403.5	L7.5	T0	L6.5+T2	$\Delta F110W = 0.526 \pm 0.015$	$\Delta F170M = 0.820 \pm 0.013$	1
2MASS J05185995-2828372	L7.5p	T1p	L6:+T4:	$\Delta F110W = 0.8 \pm 0.5$	$\Delta F170M = 0.9 \pm 0.6$	2,3
SDSS J102109.69-030420.1	T4 $\pm$ 2	T3	T1+T5	$\Delta F110W = 0.06 \pm 0.04$	$\Delta F170M = 1.030 \pm 0.019$	3
2MASS J14044941-3159329	T0	T2.5	T1+T5	$\Delta J = -0.53 \pm 0.16$	$\Delta K = 1.20 \pm 0.21$	4
SDSS J153417.05+161546.1AB	...	T3.5	T1.5+T5.5	$\Delta J = -0.17 \pm 0.04$	$\Delta K = 1.07 \pm 0.05$	5
DENIS-P J225210.73-173013.4	...	L7.5	L6+T2	$\Delta F110W = 0.526 \pm 0.015$	$\Delta F170M = 0.820 \pm 0.013$	6

<sup>a</sup>Relative photometry defined as  $M_B - M_A$ ; negative magnitudes indicate a secondary that is brighter than the primary. Relative  $JK$  photometry listed in on the MKO system.

References. — (1) Burgasser et al. (2005b); (2) Cruz et al. (2004); (3) Burgasser et al. (2006c); (4) Looper et al. (2008a); (5) Liu et al. (2006); (6) Reid et al. (2006)

Table 4. Spectral Indices.

Index (1)	Numerator Range <sup>a</sup> (2)	Denominator Range <sup>a</sup> (3)	Feature (4)	Ref (5)
H <sub>2</sub> O-J	1.14–1.165	1.26–1.285	1.15 $\mu\text{m}$ H <sub>2</sub> O	1
CH <sub>4</sub> -J	1.315–1.34	1.26–1.285	1.32 $\mu\text{m}$ CH <sub>4</sub>	1
H <sub>2</sub> O-H	1.48–1.52	1.56–1.60	1.4 $\mu\text{m}$ H <sub>2</sub> O	1
CH <sub>4</sub> -H	1.635–1.675	1.56–1.60	1.65 $\mu\text{m}$ CH <sub>4</sub>	1
H <sub>2</sub> O-K	1.975–1.995	2.08–2.10	1.9 $\mu\text{m}$ H <sub>2</sub> O	1
CH <sub>4</sub> -K	2.215–2.255	2.08–2.12	2.2 $\mu\text{m}$ CH <sub>4</sub>	1
K/J	2.060–2.10	1.25–1.29	$J - K$ color	1
H-dip	1.61–1.64	1.56–1.59 + 1.66–1.69 <sup>b</sup>	1.65 $\mu\text{m}$ CH <sub>4</sub> <sup>c</sup>	2

<sup>a</sup>Wavelength range (in microns) over which flux density ( $f_\lambda$ ) is integrated.

<sup>b</sup>Denominator is sum of these two wavelength ranges.

<sup>c</sup>Specifically, this index samples the sharp CH<sub>4</sub> feature present in the near-infrared spectra of the suspected L dwarf plus T dwarf binaries 2MASS J0805+4812 (Burgasser 2007b) and Kelu-1A (Stumpf et al. 2008).

References. — (1) Burgasser et al. (2006b); (2) This paper.



Table 5. Binary Index Selection Criteria.<sup>a</sup>

Ordinate (1)	Abcissa (2)	Inflection Points (3)	# <sup>a</sup> (4)
H <sub>2</sub> O-J	H <sub>2</sub> O-K	(0.325,0.5), (0.65,0.7)	27
CH <sub>4</sub> -H	CH <sub>4</sub> -K	(0.6,0.35), (1,0.775)	24
CH <sub>4</sub> -H	K/J	(0.65,0.25), (1,0.375)	28
H <sub>2</sub> O-H	H-dip	(0.5,0.49), (0.875,0.49)	16
Spex SpT	H <sub>2</sub> O-J/H <sub>2</sub> O-H	(L8.5,0.925), (T1.5,0.925), (T3.5,0.85)	25
Spex SpT	H <sub>2</sub> O-J/CH <sub>4</sub> -K	(L8.5,0.625), (T4.5,0.825)	21

<sup>a</sup>See Figure 4.

<sup>b</sup>Number of sources satisfying selection criteria.

Table 6. Summary of Fits to Reference Binaries.

Source (1)	Spectral Types		$\Delta J$ (4)	$\Delta K$ (5)	$\Delta F_{110W}$ (6)	$\Delta F_{170M}$ (7)	CL <sup>a</sup> (8)
	Primary (2)	Secondary (3)					
SDSSp J042348.57-041403.5	L7.5±0.3	T2.0±0.2	0.44±0.09	1.13±0.07	0.59±0.10	0.77±0.08	>99%
2MASS J05185995-2828372	L7.5±0.2	T5.0±0.5	0.13±0.19	2.18±0.31	0.39±0.20	1.71±0.36	>99%
SDSS J102109.69-030420.1	T1.0±0.4	T5.5±0.7	0.16±0.41	1.46±0.29	0.37±0.40	1.49±0.51	>99%
2MASS J14044941-3159329	T0.0±0.2	T5.0±0.2	-0.27±0.07	1.41±0.08	-0.07±0.06	1.07±0.09	>99%
SDSS J153417.05+161546.1AB	T1.5±0.5	T5.0±1.0	0.06±0.67	1.16±0.47	0.23±0.66	1.11±0.76	>99%
DENIS-P J225210.73-173013.4	L4.5±0.7	T4.5±0.4	1.29±0.25	2.88±0.32	1.54±0.25	2.50±0.33	>99%

<sup>a</sup>Confidence limit; see Section 4.2.

Table 7. Binary Candidates.

Source (1)	Opt SpT (2)	NIR SpT (3)	SpeX SpT (4)	2MASS $J$ (5)	$J - K_s$ (6)	# Criteria (7)
Strong Candidates						
SDSS J024749.90-163112.6	...	T2 $\pm$ 1.5	T2.5 $\pm$ 1	17.19 $\pm$ 0.18	1.57 $\pm$ 0.27	6
SDSS J035104.37+481046.8	...	T1 $\pm$ 1.5	T1.5 $\pm$ 1	16.47 $\pm$ 0.13	1.47 $\pm$ 0.18	6
SDSS J103931.35+325625.5	...	T1	T1.5	16.41 $\pm$ 0.15	1.25 $\pm$ 0.22	6
2MASS J11061197+2754225	...	T2.5	T2.0	14.82 $\pm$ 0.04	1.02 $\pm$ 0.07	3
2MASS J13243559+6358284	...	T2:	T2.0 $\pm$ 1	15.60 $\pm$ 0.07	1.54 $\pm$ 0.09	3
SDSS J141530.05+572428.7	...	T3 $\pm$ 1	T3.0 $\pm$ 1	16.73 $\pm$ 0.16	<1.19	6
SDSS J143553.25+112948.6	...	T2 $\pm$ 1	T2.5 $\pm$ 1	17.14 $\pm$ 0.23	<0.23	6
SDSS J143945.86+304220.6	...	T2.5	T2.5	17.22 $\pm$ 0.23	<1.34	4
SDSS J151114.66+060742.9	...	T0 $\pm$ 2	T0.5 $\pm$ 2	16.02 $\pm$ 0.08	1.47 $\pm$ 0.13	6
SDSS J151603.03+025928.9	...	T0:	L9.5 $\pm$ 1	17.23 $\pm$ 0.20	1.80 $\pm$ 0.27	4
2MASS J1711457+223204	L6.5	...	L9.0 $\pm$ 3	17.09 $\pm$ 0.18	2.36 $\pm$ 0.20	6
2MASS J21392676+0220226	...	T1.5	T2.5 $\pm$ 1	15.26 $\pm$ 0.05	1.68 $\pm$ 0.07	3
Weak Candidates						
SDSS J011912.22+240331.6	...	T2	T2.5	17.02 $\pm$ 0.18	<-0.02	2
SDSS J075840.33+324723.4	...	T2	T2.5	14.95 $\pm$ 0.04	1.07 $\pm$ 0.07	2
SDSS J090900.73+652527.2	...	T1.5	T1.5	16.03 $\pm$ 0.09	0.86 $\pm$ 0.17	2
2MASS J09490860-1545485	...	T2	T1.5 $\pm$ 1	16.15 $\pm$ 0.12	0.92 $\pm$ 0.20	2
SDSS J120602.51+281328.7	...	T3	T3.0	16.54 $\pm$ 0.11	<0.72	2
SDSS J120747.17+024424.8	L8	T0	T0.0	15.58 $\pm$ 0.07	1.59 $\pm$ 0.09	2
SDSS J151643.01+305344.4	...	T0.5 $\pm$ 1	T1.5 $\pm$ 2	16.85 $\pm$ 0.15	1.77 $\pm$ 0.18	2
SDSS J205235.31-160929.8	...	T1 $\pm$ 1	T0.0 $\pm$ 1	16.33 $\pm$ 0.12	1.21 $\pm$ 0.19	2

Table 8. Template Fits to Strong Binary Candidates. <sup>a</sup>

Mode <sup>b</sup> (1)	Primary (2)	Pub SpT (3)	SpeX SpT (4)	Secondary (5)	Pub SpT (6)	SpeX SpT (7)	$\Delta J$ (8)	$\Delta H$ (9)	$\Delta K$ (10)	$\Delta F110W$ (11)	$\Delta F170M$ (12)	$\chi^2$ (13)
SDSS J024749.90-163112.6												
Single	1750+4222	T2.0	T1.5	...	...	...	...	...	...	...	...	12.0
L06-faint	1520+3546	T0.0	L7.5	0050-3322	T7.0	T6.5	0.75	2.07	2.52	1.06	2.75	1.99
	<Primary>	T0.0±0.2	L7.5±0.2	<Secondary>	T7.0±0.3	T6.5±0.3	0.68±0.10	1.96±0.14	2.42±0.17	0.98±0.11	2.58±0.24	...
L06-faint (IC)	1520+3546	T0.0	L7.5	1624+0029	T6.0	T6.0	0.50	1.75	2.44	0.78	2.23	1.70
	<Primary>	T0.0±0.2	L7.5±0.2	<Secondary>	T6.0±0.2	T6.0±0.2	0.50±0.04	1.74±0.05	2.42±0.07	0.77±0.04	2.21±0.08	...
L06-bright	1520+3546	T0.0	L7.5	2356-1553	T5.5	T5.5	0.36	1.47	1.97	0.61	1.87	1.81
	<Primary>	L9.5±1.1	L7.5±0.9	<Secondary>	T5.5±0.4	T5.5±0.4	0.30±0.21	1.42±0.27	2.00±0.23	0.55±0.22	1.82±0.33	...
L06-bright (IC)	1520+3546	T0.0	L7.5	1624+0029	T6.0	T6.0	0.47	1.71	2.41	0.74	2.20	1.78
	<Primary>	T0.0±0.1	L7.5±0.2	<Secondary>	T5.5±0.4	T5.5±0.3	0.43±0.05	1.58±0.11	2.14±0.19	0.69±0.06	2.00±0.15	...
SDSS J035104.37+481046.8												
Single	0858+3256	T1.0	T0.0	...	...	...	...	...	...	...	...	21.3
L06-faint	0103+1935	L6.0	L6.0	0602+4043	T4.5	T4.5	0.06	1.28	2.27	0.28	1.65	6.64
	<Primary>	L6.5±0.7	L6.0±0.7	<Secondary>	T5.0±0.7	T5.5±0.6	0.31±0.31	1.56±0.33	2.48±0.31	0.56±0.32	2.00±0.44	...
L06-faint (IC)	0318-3421	L7.0	L6.5	1546-3325	T5.5	T5.5	0.18	1.53	2.52	0.45	1.97	6.68
	<Primary>	L6.5±1.1	L6.5±0.6	<Secondary>	T5.0±0.7	T5.0±0.6	0.26±0.30	1.51±0.33	2.48±0.30	0.49±0.31	1.94±0.41	...
L06-bright	0205+1251	L5.0	L5.5	2154+5942	T5.0	T5.5	0.21	1.49	2.37	0.43	1.93	6.55
	<Primary>	L5.5±0.6	L5.5±0.5	<Secondary>	T5.0±0.5	T5.0±0.5	0.27±0.26	1.48±0.28	2.38±0.26	0.50±0.27	1.90±0.34	...
L06-bright (IC)	0205+1251	L5.0	L5.5	2124+0059	T5.0	T5.0	0.16	1.35	2.14	0.38	1.73	6.63
	<Primary>	L5.5±1.0	L5.5±0.6	<Secondary>	T5.0±0.6	T5.5±0.5	0.29±0.31	1.52±0.34	2.42±0.32	0.53±0.32	1.94±0.40	...
SDSS J103931.35+325625.5												
Single	0151+1244	T1.0	T0.0	...	...	...	...	...	...	...	...	27.1
L06-faint	1043+1213	L7.0	L8.0	2254+3123	T4.0	T4.0	0.29	1.10	1.79	0.48	1.31	2.56
	<Primary>	L7.0±0.2	L8.0±0.2	<Secondary>	T4.0±0.2	T4.0±0.2	0.26±0.09	1.08±0.09	1.78±0.08	0.46±0.09	1.29±0.09	...
L06-faint (IC)	1007+1930	L8.0	L9.5	1546+4932	T2.5	T3.0	-0.38	0.14	0.86	-0.27	0.29	3.42
	<Primary>	L8.0±0.4	L9.0±0.7	<Secondary>	T3.0±0.8	T3.0±0.5	-0.25±0.19	0.37±0.32	1.06±0.34	-0.11±0.23	0.53±0.36	...
L06-bright	0857+5708	L8.0	L8.5	1546+4932	T2.5	T3.0	-1.09	-0.38	0.49	-0.97	-0.20	3.41
	<Primary>	L7.5±0.3	L7.0±1.1	<Secondary>	T2.0±0.3	T3.0±0.2	-1.05±0.14	-0.37±0.10	0.44±0.07	-0.90±0.13	-0.22±0.09	...
L06-bright (IC)	0857+5708	L8.0	L8.5	1546+4932	T2.5	T3.0	-0.99	-0.29	0.58	-0.87	-0.11	3.24
	<Primary>	L8.0±0.4	L8.0±0.8	<Secondary>	T2.5±0.5	T3.0±0.5	-0.71±0.37	-0.06±0.35	0.69±0.28	-0.57±0.41	0.08±0.33	...
2MASS J11061197+2754225												
Single	0136+0933	T2.5	T2.0	...	...	...	...	...	...	...	...	27.9
L06-faint	1520+3546	T0.0	L7.5	0000+2554	T4.5	T4.5	-0.31	0.57	1.24	-0.10	0.82	5.18

Table 8—Continued

Mode <sup>b</sup> (1)	Primary (2)	Pub SpT (3)	SpeX SpT (4)	Secondary (5)	Pub SpT (6)	SpeX SpT (7)	$\Delta J$ (8)	$\Delta H$ (9)	$\Delta K$ (10)	$\Delta F110W$ (11)	$\Delta F170M$ (12)	$\chi^2$ (13)
L06-faint (IC)	<Primary>	T0.0±0.2	L7.5±0.2	<Secondary>	T4.5±0.2	T4.5±0.2	-0.37±0.06	0.52±0.07	1.16±0.09	-0.16±0.05	0.76±0.11	...
	2043-1551	L9.0	L9.0	0559-1404	T4.5	T4.5	-0.18	0.78	1.42	0.02	1.08	7.13
L06-bright	<Primary>	L9.0±0.2	L9.0±0.2	<Secondary>	T4.5±0.2	T4.5±0.2	-0.19±0.03	0.77±0.05	1.43±0.04	0.01±0.04	1.08±0.05	...
	1520+3546	T0.0	L7.5	0559-1404	T4.5	T4.5	-0.34	0.64	1.30	-0.13	0.95	4.74
L06-bright (IC)	<Primary>	T0.0±0.2	L7.5±0.2	<Secondary>	T4.5±0.2	T4.5±0.2	-0.36±0.05	0.60±0.08	1.26±0.10	-0.15±0.04	0.90±0.13	...
	1520+3546	T0.0	L7.5	0559-1404	T4.5	T4.5	-0.22	0.76	1.42	-0.01	1.07	5.98
	<Primary>	L9.5±0.4	L8.0±0.8	<Secondary>	T4.5±0.2	T4.5±0.3	-0.26±0.11	0.69±0.18	1.35±0.16	-0.06±0.13	0.99±0.21	...
2MASS J13243559+6358284												
Single	0830+4828	L9.0	L9.5	...	...	...	...	...	...	...	...	32.0
L06-faint	1043+2225	L8.0	L9.0	1214+6316	T3.5	T4.0	-0.10	0.63	1.36	0.07	0.78	6.17
	<Primary>	L8.0±0.2	L9.5±0.2	<Secondary>	T3.5±0.2	T4.0±0.2	-0.05±0.06	0.65±0.03	1.36±0.01	0.13±0.07	0.80±0.02	...
L06-faint (IC)	1043+2225	L8.0	L9.0	1214+6316	T3.5	T4.0	-0.21	0.53	1.25	-0.03	0.67	6.03
	<Primary>	L8.0±0.2	L9.0±0.2	<Secondary>	T3.5±0.2	T4.0±0.2	-0.20±0.03	0.53±0.02	1.25±0.01	-0.03±0.03	0.67±0.01	...
L06-bright	0857+5708	L8.0	L8.5	1214+6316	T3.5	T4.0	-0.80	0.03	0.86	-0.61	0.19	8.52
	<Primary>	L8.0±0.4	L8.5±1.0	<Secondary>	T3.5±0.9	T4.0±0.9	-0.63±0.41	0.15±0.52	0.91±0.50	-0.45±0.41	0.32±0.61	...
L06-bright (IC)	1043+2225	L8.0	L9.0	1214+6316	T3.5	T4.0	-0.38	0.35	1.07	-0.21	0.50	7.12
	<Primary>	L8.0±0.2	L9.0±0.5	<Secondary>	T3.5±0.2	T4.0±0.2	-0.46±0.10	0.32±0.06	1.09±0.06	-0.28±0.09	0.47±0.06	...
SDSS J141530.05+572428.7												
Single	1254-0122	T2.0	T2.0	...	...	...	...	...	...	...	...	11.1
L06-faint	1523+3014	L8.0	L7.5	0741+2351	T5.0	T5.5	-0.22	1.06	1.91	0.03	1.47	1.73
	<Primary>	L8.0±0.5	L8.5±1.0	<Secondary>	T5.0±0.3	T5.5±0.2	-0.18±0.12	1.07±0.11	1.93±0.09	0.05±0.13	1.49±0.13	...
L06-faint (IC)	1219+3128	L8.0	L9.5	1624+0029	T6.0	T6.0	-0.18	1.13	2.02	0.06	1.65	1.59
	<Primary>	L8.5±0.8	L9.5±0.2	<Secondary>	T5.5±0.4	T5.5±0.3	-0.28±0.13	0.97±0.14	1.85±0.15	-0.06±0.15	1.43±0.18	...
L06-bright	1523+3014	L8.0	L7.5	0741+2351	T5.0	T5.5	-0.41	0.87	1.72	-0.16	1.28	1.38
	<Primary>	L8.0±0.5	L7.5±0.8	<Secondary>	T5.0±0.2	T5.5±0.2	-0.46±0.13	0.83±0.09	1.75±0.05	-0.22±0.13	1.26±0.09	...
L06-bright (IC)	1219+3128	L8.0	L9.5	0741+2351	T5.0	T5.5	-0.16	1.06	1.93	0.04	1.49	1.82
	<Primary>	L8.5±1.0	L9.0±0.6	<Secondary>	T5.0±0.3	T5.5±0.3	-0.26±0.25	0.96±0.22	1.84±0.18	-0.05±0.27	1.37±0.24	...
SDSS J143553.25+112948.6												
Single	1209-1004	T3.0	T3.0	...	...	...	...	...	...	...	...	42.4
L06-faint	1043+1213	L7.0	L8.0	1615+1340	T6.0	T6.0	0.48	1.89	2.66	0.71	2.44	19.9
	<Primary>	L7.5±0.4	L8.0±0.3	<Secondary>	T6.0±0.3	T6.0±0.2	0.41±0.12	1.75±0.23	2.51±0.25	0.65±0.10	2.26±0.30	...
L06-faint (IC)	1520+3546	T0.0	L7.5	1615+1340	T6.0	T6.0	0.48	1.82	2.47	0.73	2.36	15.9
	<Primary>	T0.0±0.6	L7.5±0.2	<Secondary>	T6.0±0.3	T6.0±0.2	0.48±0.04	1.80±0.07	2.45±0.08	0.72±0.04	2.33±0.09	...

Table 8—Continued

Mode <sup>b</sup> (1)	Primary (2)	Pub SpT (3)	SpeX SpT (4)	Secondary (5)	Pub SpT (6)	SpeX SpT (7)	$\Delta J$ (8)	$\Delta H$ (9)	$\Delta K$ (10)	$\Delta F110W$ (11)	$\Delta F170M$ (12)	$\chi^2$ (13)
L06-bright	1520+3546	T0.0	L7.5	1615+1340	T6.0	T6.0	0.36	1.70	2.35	0.61	2.24	16.6
	<Primary>	T0.0±0.2	L7.5±0.2	<Secondary>	T6.0±0.2	T6.0±0.2	0.36±0.02	1.70±0.04	2.35±0.03	0.61±0.02	2.24±0.05	...
L06-bright (IC)	1520+3546	T0.0	L7.5	1615+1340	T6.0	T6.0	0.46	1.80	2.44	0.71	2.34	14.8
	<Primary>	L9.5±0.7	L7.5±0.2	<Secondary>	T6.0±0.4	T6.0±0.2	0.45±0.02	1.77±0.06	2.41±0.08	0.70±0.02	2.30±0.09	...
SDSS J143945.86+304220.6												
Single	0136+0933	T2.5	T2.0	...	...	...	...	...	...	...	...	7.03
L06-faint	0151+1244	T1.0	T0.0	2356-1553	T5.5	T5.5	0.26	1.20	1.44	0.48	1.56	4.85
	<Primary>	T1.0±0.2	T0.0±0.3	<Secondary>	T5.0±0.6	T5.0±0.5	0.10±0.17	0.95±0.26	1.21±0.23	0.31±0.18	1.24±0.33	...
L06-faint (IC)	0151+1244	T1.0	T0.0	2151-4853	T4.0	T4.0	0.03	0.74	1.10	0.23	0.90	4.37
	<Primary>	T1.0±0.2	T0.0±0.2	<Secondary>	T4.5±0.5	T5.0±0.5	0.21±0.15	1.06±0.26	1.35±0.21	0.41±0.16	1.31±0.33	...
L06-bright	0151+1244	T1.0	T0.0	2151-4853	T4.0	T4.0	-0.14	0.57	0.93	0.06	0.73	4.59
	<Primary>	T1.0±0.2	T0.0±0.2	<Secondary>	T5.0±0.6	T5.0±0.6	0.24±0.30	1.11±0.40	1.44±0.39	0.45±0.31	1.38±0.49	...
L06-bright (IC)	0151+1244	T1.0	T0.0	2151-4853	T4.0	T4.0	0.06	0.77	1.13	0.26	0.93	4.45
	<Primary>	T1.0±0.2	T0.0±0.3	<Secondary>	T4.5±0.7	T5.0±0.6	0.27±0.31	1.12±0.41	1.49±0.41	0.48±0.31	1.37±0.51	...
SDSS J151114.66+060742.9												
Single	1030+0213	L9.5	L9.0	...	...	...	...	...	...	...	...	5.91
L06-faint	0624-4521	L5.0	L5.0	0741+2351	T5.0	T5.5	0.62	1.90	2.80	0.85	2.33	1.11
	<Primary>	L5.5±0.8	L5.0±0.3	<Secondary>	T5.0±0.4	T5.0±0.4	0.55±0.26	1.78±0.32	2.66±0.32	0.79±0.26	2.20±0.37	...
L06-faint (IC)	0624-4521	L5.0	L5.0	0741+2351	T5.0	T5.5	0.78	2.06	2.96	1.01	2.49	1.29
	<Primary>	L5.5±0.7	L5.0±0.4	<Secondary>	T5.5±0.7	T5.5±0.6	0.71±0.24	1.97±0.30	2.94±0.34	0.95±0.26	2.43±0.37	...
L06-bright	0624-4521	L5.0	L5.0	0516-0445	T5.5	T5.5	0.68	1.93	2.72	0.94	2.41	1.08
	<Primary>	L5.0±0.3	L5.0±0.4	<Secondary>	T5.5±0.4	T5.5±0.4	0.57±0.20	1.81±0.25	2.67±0.26	0.81±0.21	2.25±0.31	...
L06-bright (IC)	0624-4521	L5.0	L5.0	1624+0029	T6.0	T6.0	0.71	2.08	2.99	0.97	2.60	1.08
	<Primary>	L5.0±0.5	L5.0±0.3	<Secondary>	T5.5±0.6	T5.5±0.5	0.60±0.18	1.86±0.25	2.72±0.29	0.84±0.18	2.30±0.33	...
SDSS J151603.03+025928.9												
Single	0328+2302	L9.5	L9.0	...	...	...	...	...	...	...	...	2.85
L06-faint	0318-3421	L7.0	L6.5	1048+0919	T2.5	T2.5	-0.05	0.57	1.33	0.10	0.72	1.97
	<Primary>	L7.5±1.1	L6.5±1.1	<Secondary>	T2.5±2.2	T2.5±2.4	0.30±0.65	0.87±0.90	1.41±1.02	0.42±0.71	1.08±1.17	...
L06-faint (IC)	0318-3421	L7.0	L6.5	1048+0919	T2.5	T2.5	0.07	0.69	1.45	0.22	0.84	1.98
	<Primary>	L7.5±0.9	L6.5±0.8	<Secondary>	T3.0±2.3	T3.0±2.5	0.48±0.67	1.13±0.96	1.74±1.06	0.63±0.73	1.38±1.25	...
L06-bright	1044+0429	L7.0	L7.0	1217-0311	T7.5	T7.5	1.85	3.18	3.71	2.18	4.02	1.92
	<Primary>	L7.5±1.3	L7.0±0.8	<Secondary>	T4.0±3.5	T3.5±3.9	0.89±0.90	1.52±1.48	1.81±1.80	1.03±1.04	1.93±1.91	...
L06-bright (IC)	1044+0429	L7.0	L7.0	0039+2115	T7.5	T8.0	1.82	3.19	3.94	2.11	4.07	1.90

Table 8—Continued

Mode <sup>b</sup> (1)	Primary (2)	Pub SpT (3)	SpeX SpT (4)	Secondary (5)	Pub SpT (6)	SpeX SpT (7)	$\Delta J$ (8)	$\Delta H$ (9)	$\Delta K$ (10)	$\Delta F110W$ (11)	$\Delta F170M$ (12)	$\chi^2$ (13)
	<Primary>	L7.5±1.1	L7.0±1.0	<Secondary>	T4.5±3.3	T4.5±3.7	1.00±0.90	1.80±1.46	2.24±1.76	1.18±1.03	2.29±1.89	...
2MASS J1711457+223204												
Single	0858+3256	T1.0	T0.0	...	...	...	...	...	...	...	...	3.44
L06-faint	0624-4521	L5.0	L5.0	1828-4849	T5.5	T5.5	1.01	2.21	3.02	1.23	2.64	1.49
	<Primary>	L5.0±0.4	L5.5±0.5	<Secondary>	T5.5±1.2	T5.5±1.1	0.92±0.30	2.10±0.48	3.05±0.60	1.15±0.34	2.57±0.64	...
L06-faint (IC)	0624-4521	L5.0	L5.0	1346-0031	T6.5	T6.0	1.20	2.59	3.38	1.47	3.17	1.49
	<Primary>	L5.0±0.8	L5.5±0.7	<Secondary>	T5.5±1.4	T6.0±1.3	1.02±0.42	2.22±0.60	3.09±0.67	1.26±0.46	2.73±0.81	...
L06-bright	0624-4521	L5.0	L5.0	1114-2618	T7.5	T7.5	1.29	2.65	4.24	1.58	3.48	1.56
	<Primary>	L5.0±0.7	L5.5±0.6	<Secondary>	T6.5±0.8	T6.5±0.8	1.19±0.26	2.51±0.31	3.52±0.49	1.46±0.27	3.13±0.45	...
L06-bright (IC)	0624-4521	L5.0	L5.0	1346-0031	T6.5	T6.0	1.07	2.45	3.25	1.34	3.04	1.49
	<Primary>	L4.5±0.8	L5.5±0.8	<Secondary>	T6.5±1.1	T6.5±1.1	1.20±0.38	2.47±0.51	3.36±0.65	1.46±0.40	3.11±0.73	...
2MASS J21392676+0220226												
Single	1750+4222	T2.0	T1.5	...	...	...	...	...	...	...	...	3.29
L06-faint	0830+4828	L9.0	L9.5	1214+6316	T3.5	T4.0	-0.15	0.50	1.12	0.00	0.63	1.83
	<Primary>	L8.5±0.7	L9.0±0.6	<Secondary>	T3.5±1.0	T4.0±1.0	-0.14±0.21	0.62±0.38	1.28±0.45	0.02±0.24	0.82±0.47	...
L06-faint (IC)	0830+4828	L9.0	L9.5	1214+6316	T3.5	T4.0	-0.10	0.55	1.16	0.05	0.67	1.84
	<Primary>	L8.5±0.7	L9.5±0.4	<Secondary>	T3.5±1.1	T3.5±1.1	-0.21±0.27	0.51±0.38	1.15±0.41	-0.06±0.31	0.69±0.45	...
L06-bright	0028+2249	L7.0	L5.0	1214+6316	T3.5	T4.0	-0.38	0.27	0.98	-0.16	0.42	1.71
	<Primary>	L8.0±0.8	L8.0±1.6	<Secondary>	T3.5±0.8	T3.5±0.9	-0.46±0.26	0.19±0.34	0.84±0.37	-0.28±0.29	0.33±0.39	...
L06-bright (IC)	0830+4828	L9.0	L9.5	1214+6316	T3.5	T4.0	-0.20	0.45	1.06	-0.06	0.57	1.82
	<Primary>	L8.5±1.0	L9.0±0.8	<Secondary>	T3.0±1.1	T3.5±1.2	-0.38±0.35	0.29±0.49	0.91±0.54	-0.23±0.39	0.44±0.57	...

<sup>a</sup>For each source, we list the parameters of the best-fitting template (minimum  $\chi^2$ ) for each of the fitting modes; for the composite fits, we also give mean parameters for the primary (<Primary>) and secondary (<Secondary>), as well as the predicted relative magnitudes in the MKO *JHK* and HST/NICMOS *F110W* and *F170M* bands.

<sup>b</sup>Spectral fitting mode: Single: fits to single templates; L06-faint and L06-bright: fits to composite templates using the Liu et al. (2006) faint and bright  $M_K$ /spectral type relations, respectively; L06-faint (IC) and L06-bright (IC): fits to composite templates using the Liu et al. (2006) faint and bright relations, respectively, and the index-based spectral classifications.

Table 9. Template Fits to Weak Binary Candidates. <sup>a</sup>

Mode <sup>b</sup> (1)	Primary (2)	Pub SpT (3)	SpeX SpT (4)	Secondary (5)	Pub SpT (6)	SpeX SpT (7)	$\Delta J$ (8)	$\Delta H$ (9)	$\Delta K$ (10)	$\Delta F110W$ (11)	$\Delta F170M$ (12)	$\chi^2$ (13)
SDSS J011912.22+240331.6												
Single	1254-0122	T2.0	T2.0	...	...	...	...	...	...	...	...	25.4
L06-faint	0858+3256	T1.0	T0.0	2254+3123	T4.0	T4.0	-0.60	0.08	0.83	-0.38	0.29	15.7
	<Primary>	T0.0±0.7	L9.0±0.9	<Secondary>	T4.0±0.4	T4.5±0.5	-0.42±0.19	0.43±0.33	1.12±0.28	-0.21±0.20	0.66±0.37	...
L06-faint (IC)	0858+3256	T1.0	T0.0	1750+1759	T3.5	T3.0	-0.69	-0.06	0.79	-0.55	0.14	14.9
	<Primary>	T0.0±1.0	T0.0±0.4	<Secondary>	T4.0±0.5	T3.5±0.7	-0.54±0.23	0.20±0.39	0.99±0.33	-0.39±0.27	0.42±0.45	...
L06-bright	0328+2302	L9.5	L9.0	2151-4853	T4.0	T4.0	-0.47	0.38	1.04	-0.26	0.58	14.2
	<Primary>	L9.0±0.7	L9.0±0.8	<Secondary>	T4.0±0.3	T4.0±0.5	-0.54±0.11	0.25±0.20	0.97±0.16	-0.36±0.16	0.46±0.21	...
L06-bright (IC)	1030+0213	L9.5	L9.0	1750+1759	T3.5	T3.0	-0.71	-0.05	0.70	-0.59	0.16	13.5
	<Primary>	L9.0±0.7	L9.0±0.4	<Secondary>	T3.5±0.2	T3.5±0.3	-0.75±0.16	-0.05±0.19	0.73±0.15	-0.63±0.19	0.17±0.19	...
SDSS J075840.33+324723.4												
Single	1254-0122	T2.0	T2.0	...	...	...	...	...	...	...	...	34.8
L06-faint	2047-0718	T0.0	L7.5	1214+6316	T3.5	T4.0	-0.13	0.43	0.89	0.08	0.51	47.1
	<Primary>	T0.0±0.8	L9.0±1.1	<Secondary>	T3.5±0.8	T4.0±0.6	-0.31±0.25	0.32±0.37	0.93±0.41	-0.12±0.26	0.46±0.49	...
L06-faint (IC)	0151+1244	T1.0	T0.0	1214+6316	T3.5	T4.0	0.40	0.82	1.04	0.59	0.87	49.4
	<Primary>	L9.5±0.9	T0.0±0.2	<Secondary>	T4.0±0.4	T4.0±0.3	-0.12±0.36	0.59±0.17	1.18±0.14	0.04±0.38	0.78±0.12	...
L06-bright	2047-0718	T0.0	L7.5	1214+6316	T3.5	T4.0	-0.25	0.31	0.77	-0.04	0.39	36.7
	<Primary>	L9.5±0.6	L8.5±0.8	<Secondary>	T3.5±0.3	T4.0±0.3	-0.28±0.11	0.26±0.14	0.77±0.12	-0.08±0.13	0.35±0.14	...
L06-bright (IC)	0151+1244	T1.0	T0.0	1214+6316	T3.5	T4.0	0.40	0.81	1.03	0.58	0.87	49.4
	<Primary>	L9.5±0.9	T0.0±0.3	<Secondary>	T4.0±0.5	T4.0±0.4	-0.14±0.40	0.57±0.28	1.16±0.24	0.03±0.41	0.76±0.30	...
SDSS J090900.73+652527.2												
Single	0136+0933	T2.5	T2.0	...	...	...	...	...	...	...	...	4.76
L06-faint	1750+4222	T2.0	T1.5	1048+0919	T2.5	T2.5	-0.16	-0.07	0.13	-0.16	0.00	2.53
	<Primary>	T1.5±0.5	T0.5±0.7	<Secondary>	T2.5±0.3	T2.5±0.5	-0.12±0.10	0.10±0.17	0.30±0.19	-0.07±0.11	0.17±0.18	...
L06-faint (IC)	1750+4222	T2.0	T1.5	1048+0919	T2.5	T2.5	-0.06	0.03	0.24	-0.05	0.10	2.52
	<Primary>	T1.5±0.5	T1.0±0.6	<Secondary>	T2.5±0.3	T2.5±0.4	0.02±0.10	0.22±0.22	0.42±0.25	0.07±0.14	0.29±0.24	...
L06-bright	1750+4222	T2.0	T1.5	1048+0919	T2.5	T2.5	-0.16	-0.07	0.13	-0.15	0.00	2.53
	<Primary>	T1.0±0.8	T0.5±1.1	<Secondary>	T3.0±0.4	T2.5±0.5	-0.20±0.17	0.08±0.14	0.34±0.23	-0.13±0.15	0.17±0.16	...
L06-bright (IC)	1750+4222	T2.0	T1.5	1048+0919	T2.5	T2.5	-0.09	0.00	0.20	-0.08	0.07	2.52
	<Primary>	T1.5±0.8	T0.5±1.0	<Secondary>	T2.5±0.4	T2.5±0.5	-0.12±0.18	0.15±0.17	0.40±0.22	-0.06±0.17	0.23±0.18	...
2MASS J09490860-1545485												
Single	0136+0933	T2.5	T2.0	...	...	...	...	...	...	...	...	27.2
L06-faint	1632+4150	T1.0	T1.0	1122-3512	T2.0	T2.0	-0.09	0.09	0.24	-0.07	0.15	13.7

Table 9—Continued

Mode <sup>b</sup> (1)	Primary (2)	Pub SpT (3)	SpeX SpT (4)	Secondary (5)	Pub SpT (6)	SpeX SpT (7)	$\Delta J$ (8)	$\Delta H$ (9)	$\Delta K$ (10)	$\Delta F110W$ (11)	$\Delta F170M$ (12)	$\chi^2$ (13)
L06-faint (IC)	<Primary>	T1.0±0.2	T1.0±0.2	<Secondary>	T2.0±0.2	T2.0±0.3	-0.07±0.05	0.12±0.06	0.26±0.05	-0.04±0.06	0.18±0.06	...
	1632+4150	T1.0	T1.0	1122-3512	T2.0	T2.0	-0.09	0.09	0.24	-0.07	0.15	13.7
L06-bright	<Primary>	T1.0±0.2	T1.0±0.2	<Secondary>	T2.0±0.2	T2.0±0.3	-0.05±0.09	0.13±0.10	0.28±0.09	-0.02±0.10	0.20±0.10	...
	1632+4150	T1.0	T1.0	1122-3512	T2.0	T2.0	-0.16	0.02	0.17	-0.14	0.08	13.7
L06-bright (IC)	<Primary>	T1.0±0.2	T1.0±0.2	<Secondary>	T2.0±0.4	T2.0±0.4	-0.13±0.14	0.06±0.20	0.21±0.19	-0.10±0.15	0.13±0.23	...
	1632+4150	T1.0	T1.0	1122-3512	T2.0	T2.0	-0.15	0.03	0.18	-0.13	0.09	13.7
	<Primary>	T1.0±0.2	T1.0±0.2	<Secondary>	T2.0±0.3	T2.0±0.4	-0.10±0.12	0.08±0.14	0.23±0.13	-0.08±0.13	0.15±0.16	...
SDSS J120602.51+281328.7												
Single	1750+1759	T3.5	T3.0	...	...	...	...	...	...	...	...	27.3
L06-faint	1122-3512	T2.0	T2.0	1901+4718	T5.0	T5.0	0.23	0.84	0.98	0.31	1.05	22.8
L06-faint (IC)	<Primary>	T2.0±0.2	T2.0±0.2	<Secondary>	T4.5±0.4	T4.5±0.3	0.16±0.10	0.73±0.19	0.86±0.18	0.24±0.11	0.93±0.22	...
	1122-3512	T2.0	T2.0	1901+4718	T5.0	T5.0	0.14	0.75	0.88	0.21	0.96	23.2
L06-bright	<Primary>	T2.0±0.3	T2.0±0.3	<Secondary>	T4.5±0.5	T4.5±0.5	0.08±0.14	0.62±0.27	0.75±0.26	0.16±0.16	0.81±0.32	...
	1122-3512	T2.0	T2.0	0742+2055	T5.0	T5.0	0.49	1.17	1.37	0.61	1.44	26.7
L06-bright (IC)	<Primary>	T2.0±0.3	T2.0±0.3	<Secondary>	T5.0±0.5	T5.0±0.6	0.42±0.19	1.03±0.34	1.20±0.39	0.52±0.22	1.26±0.40	...
	1122-3512	T2.0	T2.0	1901+4718	T5.0	T5.0	0.48	1.10	1.23	0.56	1.31	25.7
	<Primary>	T2.0±0.4	T2.5±0.4	<Secondary>	T4.5±0.6	T4.5±0.7	0.37±0.33	0.91±0.52	1.06±0.51	0.45±0.37	1.11±0.59	...
SDSS J120747.17+024424.8												
Single	0151+1244	T1.0	T0.0	...	...	...	...	...	...	...	...	44.6
L06-faint	1036-3441	L6.0	L6.5	1209-1004	T3.0	T3.0	0.74	1.24	1.77	0.87	1.38	6.14
L06-faint (IC)	<Primary>	L6.5±0.7	L6.5±0.4	<Secondary>	T2.5±0.5	T2.5±0.5	0.48±0.28	0.90±0.31	1.45±0.28	0.62±0.30	1.01±0.31	...
	1036-3441	L6.0	L6.5	1048+0919	T2.5	T2.5	0.47	0.87	1.36	0.59	0.97	4.95
L06-bright	<Primary>	L6.0±0.5	L7.0±0.2	<Secondary>	T2.5±0.3	T2.0±0.2	0.43±0.13	0.83±0.12	1.32±0.06	0.55±0.13	0.92±0.11	...
	1515+4847	L6.0	L5.5	1254-0122	T2.0	T2.0	0.11	0.34	0.68	0.26	0.37	4.38
L06-bright (IC)	<Primary>	L5.5±0.4	L5.5±0.2	<Secondary>	T2.0±0.2	T2.0±0.2	0.20±0.16	0.43±0.15	0.76±0.14	0.36±0.17	0.46±0.15	...
	1515+4847	L6.0	L5.5	1254-0122	T2.0	T2.0	0.24	0.47	0.82	0.40	0.51	4.18
	<Primary>	L6.0±0.3	L5.5±0.2	<Secondary>	T2.0±0.2	T2.0±0.2	0.24±0.01	0.47±0.02	0.81±0.02	0.40±0.00	0.50±0.02	...
SDSS J151643.01+305344.4												
Single	1007+1930	L8.0	L9.5	...	...	...	...	...	...	...	...	8.38
L06-faint	1043+2225	L8.0	L9.0	0830+4828	L9.0	L9.5	0.05	0.13	0.24	0.07	0.16	7.28
L06-faint (IC)	<Primary>	L8.0±0.2	L9.0±0.5	<Secondary>	L9.5±1.2	T0.0±0.9	0.06±0.24	0.21±0.33	0.34±0.39	0.08±0.25	0.25±0.42	...
	1043+2225	L8.0	L9.0	0830+4828	L9.0	L9.5	-0.10	-0.02	0.09	-0.08	0.00	7.22
	<Primary>	L8.0±0.2	L9.0±0.6	<Secondary>	L9.5±0.9	T0.0±0.8	-0.09±0.13	0.05±0.23	0.19±0.35	-0.07±0.16	0.08±0.27	...



Table 9—Continued

Mode <sup>b</sup> (1)	Primary (2)	Pub SpT (3)	SpeX SpT (4)	Secondary (5)	Pub SpT (6)	SpeX SpT (7)	$\Delta J$ (8)	$\Delta H$ (9)	$\Delta K$ (10)	$\Delta F110W$ (11)	$\Delta F170M$ (12)	$\chi^2$ (13)
L06-bright	1043+2225	L8.0	L9.0	0830+4828	L9.0	L9.5	-0.14	-0.06	0.06	-0.11	-0.03	7.20
	<Primary>	L8.0±0.2	L9.0±0.5	<Secondary>	T0.0±2.8	T0.5±2.6	-0.01±0.55	0.29±0.97	0.50±1.18	0.04±0.63	0.42±1.25	...
L06-bright (IC)	1043+2225	L8.0	L9.0	0830+4828	L9.0	L9.5	-0.18	-0.10	0.02	-0.15	-0.07	7.19
	<Primary>	L8.0±0.2	L9.0±0.5	<Secondary>	T0.0±2.7	T0.5±2.6	-0.06±0.51	0.24±0.94	0.46±1.17	-0.01±0.60	0.37±1.22	...
SDSS J205235.31-160929.8												
Single	0151+1244	T1.0	T0.0	...	...	...	...	...	...	...	...	6.23
L06-faint	1155+0559	L7.5	L7.0	1122-3512	T2.0	T2.0	-0.01	0.40	1.07	0.11	0.53	2.59
	<Primary>	L7.5±0.6	L7.0±0.5	<Secondary>	T2.0±0.2	T2.0±0.2	0.04±0.18	0.45±0.18	1.10±0.17	0.16±0.18	0.58±0.18	...
L06-faint (IC)	2047-0718	T0.0	L7.5	1048+0919	T2.5	T2.5	0.29	0.70	1.13	0.41	0.78	2.19
	<Primary>	L9.5±1.3	L7.5±0.3	<Secondary>	T2.5±0.3	T2.0±0.2	0.29±0.06	0.70±0.06	1.13±0.07	0.41±0.05	0.77±0.05	...
L06-bright	1036-3441	L6.0	L6.5	0136+0933	T2.5	T2.0	-0.01	0.39	0.81	0.11	0.46	2.94
	<Primary>	L6.0±0.6	L6.5±0.4	<Secondary>	T2.5±0.2	T2.0±0.2	-0.08±0.16	0.32±0.15	0.78±0.13	0.05±0.16	0.40±0.15	...
L06-bright (IC)	1526+2043	L7.0	L5.5	1048+0919	T2.5	T2.5	-0.10	0.29	0.85	0.06	0.40	3.30
	<Primary>	L7.0±1.2	L6.5±0.9	<Secondary>	T2.0±0.5	T2.0±0.5	-0.17±0.26	0.18±0.24	0.69±0.25	-0.02±0.28	0.27±0.24	...

<sup>a</sup>For each source, we list the parameters of the best-fitting template (minimum  $\chi^2$ ) for each of the fitting modes; for the composite fits, we also give mean parameters for the primary (<Primary>) and secondary (<Secondary>), as well as the predicted relative magnitudes in the MKO *JHK* and HST/NICMOS *F110W* and *F170M* bands.

<sup>b</sup>Spectral fitting mode: Single: fits to single templates; L06-faint and L06-bright: fits to composite templates using the Liu et al. (2006) faint and bright  $M_K$ /spectral type relations, respectively; L06-faint (IC) and L06-bright (IC): fits to composite templates using the Liu et al. (2006) faint and bright relations, respectively, and the index-based spectral classifications.

Table 10. Summary of Binary Candidates.

Source (1)	Spectral Types				$\Delta J$ (6)	CL <sup>a</sup> (7)	AO/ HST? (8)	Notes (9)
	Pub (2)	SpeX (3)	Primary (4)	Secondary (5)				
Probable Binaries								
SDSS J011912.22+240331.6	T2	T2.5	T0.0±0.7	T4.0±0.4	-0.42±0.19	>99%	No	Large $J$ flux reversal Cloudy primary?
SDSS J024749.90-163112.6	T2±1.5	T2.5±1	T0.0±0.2	T7.0±0.3	0.68±0.10	>99%	No	Old/metal-poor?
SDSS J035104.37+481046.8	T1±1.5	T1.5±1	L6.5±0.7	T5.0±0.7	0.31±0.31	>99%	No	Small $J$ flux reversal Cloudy primary?
SDSS J090900.73+652527.2	T1.5	T1.5	T1.5±0.5	T2.5±0.3	-0.12±0.10	>99%	No	Small $J$ flux reversal
2MASS J09490860-1545485	T2	T1.5±1	T1.0±0.2	T2.0±0.2	-0.07±0.05	>99%	No	Small $J$ flux reversal
SDSS J103931.35+325625.5	T1	T1.5	L7.0±0.2	T4.0±0.2	0.26±0.09	>99%	No	
2MASS J11061197+2754225	T2.5	T2.0	T0.0±0.2	T4.5±0.2	-0.37±0.06	>99%	Yes	Large $J$ flux reversal Old/metal-poor?
SDSS J120747.17+024424.8	T0	T0.0	L6.5±0.7	T2.5±0.5	0.48±0.28	>99%	No	
2MASS J13243559+6358284	T2p	T2.0±1	L8.0±0.2	T3.5±0.2	-0.05±0.06	>99%	No	Large $J$ flux reversal Cloudy primary?
SDSS J141530.05+572428.7	T3±1	T3.0±1	L8.0±0.5	T5.0±0.3	-0.18±0.12	>99%	No	Large $J$ flux reversal
SDSS J143553.25+112948.6	T2±1	T2.5±1	L7.5±0.4	T6.0±0.3	0.41±0.12	>99%	No	
SDSS J143945.86+304220.6	T2.5	T2.5	T1.0±0.2	T5.0±0.6	0.10±0.17	>99%	No	
SDSS J151114.66+060742.9	T0±2	T0.5±2	L5.5±0.8	T5.0±0.4	0.55±0.26	>99%	No	Cloudy primary?
SDSS J151603.03+025928.9	T0:	L9.5±1	L7.5±1.1	T2.5±2.2	0.30±0.65	>99%	No	Small $J$ flux reversal Cloudy primary?
2MASSI J1711457+223204	L6.5	L9.0±3	L5.0±0.4	T5.5±1.2	0.92±0.30	>99%	No <sup>b</sup>	$\pi$ measurement
SDSS J205235.31-160929.8	T1±1	T0.0±1	L7.5±0.6	T2.0±0.2	0.04±0.18	>99%	No	Small $J$ flux reversal
2MASS J21392676+0220226	T1.5	T2.0±1	L8.5±0.7	T3.5±1.0	-0.14±0.21	>99%	No	Large $J$ flux reversal
Other Sources								
SDSS J075840.33+324723.4	T2	T2.5	T0.0±0.8	T3.5±0.8	-0.31±0.25	1%	No	Clone of SDSS J1254-0122
SDSS J120602.51+281328.7	T3	T3.0	T2.0±0.2	T4.5±0.4	0.16±0.10	92%	No	Clone of SDSS J1750+1759
SDSS J151643.01+305344.4	T0.5±1	T1.5±2	L8.0±0.2	L9.5±1.2	0.06±0.24	87%	No	Thick clouds?

<sup>a</sup>Confidence limit; see Section 4.2.

<sup>b</sup>HST/WFPC2 observations in the F1042M band by Gizis et al. (2003) did not have sufficient sensitivity to detect any putative T dwarf secondary in this system.

Table 11. Estimated Binary Fractions for SpeX Sample.

SpT (1)	Number of Sources				Binary Fraction <sup>a</sup>	
	Total (2)	Resolved (3)	Candidates (4)	Non-candidates (5)	Observed (6)	Predicted <sup>b</sup> (7)
L8–L9	12	0	0	12	<13%	33%
L9–T0	9	1	0	8	$11^{+18}_{-4}\%$	49%
T0–T1	9	3	3	3	$67^{+11}_{-17}\%$	70%
T1–T2	10	1	5	4	$60^{+13}_{-16}\%$	68%
T2–T3	17	1	7	9	$47^{+12}_{-11}\%$	45%
T3–T4	7	2	1	4	$43^{+18}_{-15}\%$	23%
T4–T5	10	1	0	9	$10^{+17}_{-3}\%$	26%
T5–T6	20	1	0	19	$5^{+10}_{-2}\%$	30%
T0–T4	43	7	16	20	$53\pm7\%$	56%

<sup>a</sup>Including resolved binaries plus binary candidates.

<sup>b</sup>Based on the simulations of Burgasser (2007a), assuming a mass function  $\Psi(M) \propto M^{-0.5}$ , a constant birthrate over  $0.01 < t < 10$  Gyr, an intrinsic binary fraction  $\epsilon_b = 15\%$ , a mass ratio distribution  $P(q) \propto q^4$ , evolutionary models from Baraffe et al. (2003),  $K$ -band absolute magnitude and bolometric magnitude relations as defined in that study, and a magnitude-limited sample.

## REFERENCES

- Ackerman, A. S., & Marley, M. S. 2001, *ApJ*, 556, 872
- Allen, P. R. 2007, *ApJ*, 668, 492
- Artigau, É., Doyon, R., Lafrenière, D., Nadeau, D., Robert, J., & Albert, L. 2006, *ApJ*, 651, L57
- Baraffe, I., Chabrier, G., Barman, T. S., Allard, F., & Hauschildt, P. H. 2003, *A&A*, 402, 701
- Basri, G., & Martín, E. L. 1999, *AJ*, 118, 2460
- Basri, G., & Reiners, A. 2006, *AJ*, 132, 663
- Blake, C. H., Charbonneau, D., White, R. J., Torres, G., Marley, M. S., & Saumon, D. 2008, *ApJ*, 678, L125
- Bouy, H., Martín, E. L., Brandner, W., & Bouvier, J. 2005, *AJ*, 129, 511
- Burgasser, A. J. 2007a, *ApJ*, 659, 655
- . 2007b, *AJ*, 134, 1330
- . 2007c, *ApJ*, 658, 617
- Burgasser, A. J., Burrows, A., & Kirkpatrick, J. D. 2006a, *ApJ*, 639, 1095
- Burgasser, A. J., Geballe, T. R., Leggett, S. K., Kirkpatrick, J. D., & Golimowski, D. A. 2006b, *ApJ*, 637, 1067
- Burgasser, A. J., Kirkpatrick, J. D., Burrows, A., Liebert, J., Reid, I. N., Gizis, J. E., McGovern, M. R., Prato, L., & McLean, I. S. 2003a, *ApJ*, 592, 1186
- Burgasser, A. J., Kirkpatrick, J. D., Cruz, K. L., Reid, I. N., Leggett, S. K., Liebert, J., Burrows, A., & Brown, M. E. 2006c, *ApJS*, 166, 585
- Burgasser, A. J., Kirkpatrick, J. D., Liebert, J., & Burrows, A. 2003b, *ApJ*, 594, 510
- Burgasser, A. J., Kirkpatrick, J. D., & Lowrance, P. J. 2005a, *AJ*, 129, 2849
- Burgasser, A. J., Kirkpatrick, J. D., McElwain, M. W., Cutri, R. M., Burgasser, A. J., & Skrutskie, M. F. 2003c, *AJ*, 125, 850
- Burgasser, A. J., Liu, M. C., Ireland, M. J., Cruz, K. L., & Dupuy, T. J. 2008a, *ApJ*, 681, 579
- Burgasser, A. J.,Looper, D. L., Kirkpatrick, J. D., Cruz, K. L., & Swift, B. J. 2008b, *ApJ*, 674, 451
- Burgasser, A. J.,Looper, D. L., Kirkpatrick, J. D., & Liu, M. C. 2007a, *ApJ*, 658, 557

- Burgasser, A. J., & McElwain, M. W. 2006, *AJ*, 131, 1007
- Burgasser, A. J., McElwain, M. W., & Kirkpatrick, J. D. 2003d, *AJ*, 126, 2487
- Burgasser, A. J., McElwain, M. W., Kirkpatrick, J. D., Cruz, K. L., Tinney, C. G., & Reid, I. N. 2004, *AJ*, 127, 2856
- Burgasser, A. J., Reid, I. N., Leggett, S. K., Kirkpatrick, J. D., Liebert, J., & Burrows, A. 2005b, *ApJ*, 634, L177
- Burgasser, A. J., Reid, I. N., Siegler, N., Close, L., Allen, P., Lowrance, P., & Gizis, J. 2007b, *Protostars and Planets V*, 427
- Burgasser, A. J., et al. 1999, *ApJ*, 522, L65
- . 2000a, *AJ*, 120, 1100
- . 2000b, *ApJ*, 531, L57
- Burgasser, A. J., et al. 2002a, *ApJ*, 564, 421
- . 2002b, *ApJ*, 571, L151
- Burrows, A., Sudarsky, D., & Hubeny, I. 2006, *ApJ*, 640, 1063
- Chabrier, G., Baraffe, I., Leconte, J., Gallardo, J., & Barman, T. 2009, in *American Institute of Physics Conference Series*, Vol. 1094, American Institute of Physics Conference Series, ed. E. Stempels, 102–111
- Chiu, K., Fan, X., Leggett, S. K., Golimowski, D. A., Zheng, W., Geballe, T. R., Schneider, D. P., & Brinkmann, J. 2006, *AJ*, 131, 2722
- Cruz, K. L., Burgasser, A. J., Reid, I. N., & Liebert, J. 2004, *ApJ*, 604, L61
- Cruz, K. L., Kirkpatrick, J. D., & Burgasser, A. J. 2009, *AJ*, 137, 3345
- Cruz, K. L., et al. 2007, *AJ*, 133, 439
- Cruz, K. L., Reid, I. N., Liebert, J., Kirkpatrick, J. D., & Lowrance, P. J. 2003, *AJ*, 126, 2421
- Cushing, M. C., Marley, M. S., Saumon, D., Kelly, B. C., Vacca, W. D., Rayner, J. T., Freedman, R. S., Lodders, K., & Roellig, T. L. 2008, *ApJ*, 678, 1372
- Cushing, M. C., Rayner, J. T., & Vacca, W. D. 2005, *ApJ*, 623, 1115
- Cushing, M. C., et al. 2006, *ApJ*, 648, 614
- Cushing, M. C., Vacca, W. D., & Rayner, J. T. 2004, *PASP*, 116, 362

- Dahn, C. C., et al. 2002, *AJ*, 124, 1170
- Delfosse, X., et al. 1997, *A&A*, 327, L25
- Dupuy, T. J., Liu, M. C., & Ireland, M. J. 2009, *ApJ*, 692, 729
- Duquennoy, A., & Mayor, M. 1991, *A&A*, 248, 485
- Ellis, S. C., Tinney, C. G., Burgasser, A. J., Kirkpatrick, J. D., & McElwain, M. W. 2005, *AJ*, 130, 2347
- Epchtein, N., et al. 1997, *The Messenger*, 87, 27
- Faherty, J. K., Burgasser, A. J., Cruz, K. L., Shara, M. M., Walter, F. M., & Gelino, C. R. 2009, *AJ*, 137, 1
- Fan, X., et al. 2000, *AJ*, 119, 928
- Fischer, D. A., & Marcy, G. W. 1992, *ApJ*, 396, 178
- Geballe, T. R., et al. 2002, *ApJ*, 564, 466
- Gelino, C. R., Kulkarni, S. R., & Stephens, D. C. 2006, *PASP*, 118, 611
- Gizis, J. E. 2002, *ApJ*, 575, 484
- Gizis, J. E., Monet, D. G., Reid, I. N., Kirkpatrick, J. D., Liebert, J., & Williams, R. J. 2000, *AJ*, 120, 1085
- Gizis, J. E., Reid, I. N., Knapp, G. R., Liebert, J., Kirkpatrick, J. D., Koerner, D. W., & Burgasser, A. J. 2003, *AJ*, 125, 3302
- Goldman, B., Bouy, H., Zapatero Osorio, M. R., Stumpf, M. B., Brandner, W., & Henning, T. 2008, *A&A*, 490, 763
- Golimowski, D. A., et al. 2004, *AJ*, 127, 3516
- Hawley, S. L., et al. 2002, *AJ*, 123, 3409
- Joergens, V. 2008, *A&A*, 492, 545
- Joergens, V., & Müller, A. 2007, *ApJ*, 666, L113
- Kendall, T. R., Delfosse, X., Martín, E. L., & Forveille, T. 2004, *A&A*, 416, L17
- Kendall, T. R., Jones, H. R. A., Pinfield, D. J., Pokorny, R. S., Folkes, S., Weights, D., Jenkins, J. S., & Maun, N. 2007, *MNRAS*, 374, 445
- Kendall, T. R., Maun, N., Azzopardi, M., & Gigoyan, K. 2003, *A&A*, 403, 929

- Kirkpatrick, J. D. 2005, *ARA&A*, 43, 195
- Kirkpatrick, J. D., Reid, I. N., Liebert, J., Gizis, J. E., Burgasser, A. J., Monet, D. G., Dahn, C. C., Nelson, B., & Williams, R. J. 2000, *AJ*, 120, 447
- Kirkpatrick, J. D., et al. 1999, *ApJ*, 519, 802
- . 2008, *ApJ*, 689, 1295
- Knapp, G. R., et al. 2004, *AJ*, 127, 3553
- Leggett, S. K., Saumon, D., Marley, M. S., Geballe, T. R., Golimowski, D. A., Stephens, D., & Fan, X. 2007, *ApJ*, 655, 1079
- Leggett, S. K., et al. 2000, *ApJ*, 536, L35
- . 2002, *ApJ*, 564, 452
- Liebert, J., & Burgasser, A. J. 2007, *ApJ*, 655, 522
- Liebert, J., & Gizis, J. E. 2006, *PASP*, 118, 659
- Liebert, J., Kirkpatrick, J. D., Cruz, K. L., Reid, I. N., Burgasser, A., Tinney, C. G., & Gizis, J. E. 2003, *AJ*, 125, 343
- Liu, M. C., Dupuy, T. J., & Ireland, M. J. 2008, *ApJ*, 689, 436
- Liu, M. C., & Leggett, S. K. 2005, *ApJ*, 634, 616
- Liu, M. C., Leggett, S. K., Golimowski, D. A., Chiu, K., Fan, X., Geballe, T. R., Schneider, D. P., & Brinkmann, J. 2006, *ApJ*, 647, 1393
- Lodders, K. 2002, *ApJ*, 577, 974
- Lodieu, N., Dobbie, P. D., Deacon, N. R., Hodgkin, S. T., Hambly, N. C., & Jameson, R. F. 2007, *MNRAS*, 380, 712
- Looper, D. L., Gelino, C. R., Burgasser, A. J., & Kirkpatrick, J. D. 2008a, *ApJ*, 685, 1183
- Looper, D. L., Kirkpatrick, J. D., & Burgasser, A. J. 2007, *AJ*, 134, 1162
- Looper, D. L., et al. 2008b, *ApJ*, 686, 528
- Lucas, P. W., & Roche, P. F. 2000, *MNRAS*, 314, 858
- Luhman, K. L., et al. 2007, *ApJ*, 654, 570
- Marley, M. S., Saumon, D., Guillot, T., Freedman, R. S., Hubbard, W. B., Burrows, A., & Lunine, J. I. 1996, *Science*, 272, 1919

- Martin, E. L., Brandner, W., & Basri, G. 1999, *Science*, 283, 1718
- Martín, E. L., Delfosse, X., Basri, G., Goldman, B., Forveille, T., & Zapatero Osorio, M. R. 1999, *AJ*, 118, 2466
- McCaughrean, M. J., Close, L. M., Scholz, R.-D., Lenzen, R., Biller, B., Brandner, W., Hartung, M., & Lodieu, N. 2004, *A&A*, 413, 1029
- McElwain, M. W., & Burgasser, A. J. 2006, *AJ*, 132, 2074
- McLean, I. S., McGovern, M. R., Burgasser, A. J., Kirkpatrick, J. D., Prato, L., & Kim, S. S. 2003, *ApJ*, 596, 561
- Metchev, S. A., & Hillenbrand, L. A. 2006, *ApJ*, 651, 1166
- Metchev, S. A., Kirkpatrick, J. D., Berriman, G. B., &Looper, D. 2008, *ApJ*, 676, 1281
- Mohanty, S., Jayawardhana, R., Huélamo, N., & Mamajek, E. 2007, *ApJ*, 657, 1064
- Muench, A. A., Lada, C. J., Luhman, K. L., Muzerolle, J., & Young, E. 2007, *AJ*, 134, 411
- Mugrauer, M., Seifahrt, A., Neuhäuser, R., & Mazeh, T. 2006, *MNRAS*, 373, L31
- Nakajima, T., Tsuji, T., & Yanagisawa, K. 2004, *ApJ*, 607, 499
- Rayner, J. T., Toomey, D. W., Onaka, P. M., Denault, A. J., Stahlberger, W. E., Vacca, W. D., Cushing, M. C., & Wang, S. 2003, *PASP*, 115, 362
- Reid, I. N., Burgasser, A. J., Cruz, K. L., Kirkpatrick, J. D., & Gizis, J. E. 2001a, *AJ*, 121, 1710
- Reid, I. N., Cruz, K. L., Kirkpatrick, J. D., Allen, P. R., Mungall, F., Liebert, J., Lowrance, P., & Sweet, A. 2008, *AJ*, 136, 1290
- Reid, I. N., Gizis, J. E., Kirkpatrick, J. D., & Koerner, D. W. 2001b, *AJ*, 121, 489
- Reid, I. N., Kirkpatrick, J. D., Gizis, J. E., Dahn, C. C., Monet, D. G., Williams, R. J., Liebert, J., & Burgasser, A. J. 2000, *AJ*, 119, 369
- Reid, I. N., Lewitus, E., Burgasser, A. J., & Cruz, K. L. 2006, *ApJ*, 639, 1114
- Saumon, D., & Marley, M. S. 2008, *ApJ*, 689, 1327
- Schneider, D. P., et al. 2002, *AJ*, 123, 458
- Sheppard, S. S., & Cushing, M. C. 2009, *AJ*, 137, 304
- Siegler, N., Close, L. M., Burgasser, A. J., Cruz, K. L., Marois, C., Macintosh, B., & Barman, T. 2007, *AJ*, 133, 2320



- Simons, D. A., & Tokunaga, A. 2002, *PASP*, 114, 169
- Skrutskie, M. F., et al. 2006, *AJ*, 131, 1163
- Stassun, K. G., Mathieu, R. D., & Valenti, J. A. 2006, *Nature*, 440, 311
- Stephens, D. C., et al. 2009, *ApJ*, 702, 154
- Strauss, M. A., et al. 1999, *ApJ*, 522, L61
- Stumpf, M. B., Brandner, W., Henning, T., Bouy, H., Koehler, R., Hormuth, F., Joergens, V., & Kasper, M. 2008, *ArXiv e-prints*
- Testi, L., et al. 2001, *ApJ*, 552, L147
- Tinney, C. G., Burgasser, A. J., & Kirkpatrick, J. D. 2003, *AJ*, 126, 975
- Tinney, C. G., Burgasser, A. J., Kirkpatrick, J. D., & McElwain, M. W. 2005, *AJ*, 130, 2326
- Tokunaga, A. T., Simons, D. A., & Vacca, W. D. 2002, *PASP*, 114, 180
- Tsuji, T. 2002, *ApJ*, 575, 264
- . 2005, *ApJ*, 621, 1033
- Tsuji, T., & Nakajima, T. 2003, *ApJ*, 585, L151
- Tsuji, T., Nakajima, T., & Yanagisawa, K. 2004, *ApJ*, 607, 511
- Tsuji, T., Ohnaka, K., & Aoki, W. 1996, *A&A*, 305, L1+
- . 1999, *ApJ*, 520, L119
- Tsvetanov, Z. I., et al. 2000, *ApJ*, 531, L61
- Vacca, W. D., Cushing, M. C., & Rayner, J. T. 2003, *PASP*, 115, 389
- Vrba, F. J., et al. 2004, *AJ*, 127, 2948
- Wilson, J. C., Miller, N. A., Gizis, J. E., Skrutskie, M. F., Houck, J. R., Kirkpatrick, J. D., Burgasser, A. J., & Monet, D. G. 2003, in *IAU Symposium*, Vol. 211, *Brown Dwarfs*, ed. E. Martín, 197–+
- York, D. G., et al. 2000, *AJ*, 120, 1579
- Zapatero Osorio, M. R., Lane, B. F., Pavlenko, Y., Martín, E. L., Britton, M., & Kulkarni, S. R. 2004, *ApJ*, 615, 958